Dizziness Handicap Inventory Fact Sheet

Overview: Self-perceived questionnaire that evaluates the impacts of dizziness

Scoring: 25 questions; "yes" (4 points), "sometimes" (2 points), and "no" (3 points); total score ranges from 0-100; The higher the score the greater perceived dizziness

Domains:

- 1) Functional (9 questions, 36 points)
- 2) Emotional (9 questions, 36 points)
- 3) Physical (7 questions, 28 points)

Cut off scores:

-mild: 0-30

-moderate: 31-60

-severe: 60-100

Vestibular MDC: 17.18

Vestibular MCID: Pre-treatment and Post treatment would need to have an 18 point difference

References:

1) Jacobson, G.P., Newman, C.W. (1990). The development of the dizziness handicap inventory. Arch Otolaryngol Head Neck Surg, 116, 424-427.

2) Whitney, S.L., Wrisley, D.M., et al. (2004). "Is perception of handicap related to functional performance in persons with vestibular dysfunction?" Otol Neurotol 25(2): 139-143.



ANPT Concussion/mTBI Knowledge Translation Task force

Handout Created by:
Naseem Chatiwala, PT, DPT,
MS; Annie Fangman, PT,
DPT; Michelle Gutierrez, PT,
DSc; John Heick, PT, DPT,
PhD; Ethan Hood, PT, DPT,
MBA; Victoria Kochick, PT,
DPT; Becky Bliss, PT, DHSc,
DPT; and Pradeep Rapalli,
PT, DPT

Clinical Practice Guideline
Physical Therapy Evaluation
and Treatment After
Concussion/Mild Traumatic
Brain Injury

Journal of Orthopedic & Sports Physical Therapy; April 2020: Volume 50, Issue 4



This is for informational and educational purposes only. It does not constitute and should not be used as a substitute for medical advice, diagnosis, rehabilitation, or treatment. Patients and other members of the general public should always seek the advice of a qualified healthcare professional regarding personal health and medical conditions. The Academy of Neurologic Physical Therapy and its collaborators disclaim any liability to any party for any loss or damage by errors or omissions in this publication.

© 2023 by the Academy of Neurologic Physical Therapy.