

# FREQUENTLY ASKED QUESTIONS

## CLINICAL PRACTICE GUIDELINE TO IMPROVE LOCOMOTOR FUNCTION FOLLOWING CHRONIC STROKE, INCOMPLETE SPINAL CORD INJURY AND BRAIN INJURY



### What was the purpose of the Clinical Practice Guideline to Improve Locomotor Function Following Stroke, incomplete SCI, and Brain Injury?

The purpose of the CPG was to evaluate available evidence of the efficacy of various physical interventions to improve walking function of individuals with a history of a stroke (CVA), motor incomplete spinal cord injury (SCI), or traumatic brain injury (TBI) of > 6 months duration. [Access the publication here.](#)

### What were the populations under study?

**Individuals with a history of a stroke, motor incomplete SCI, or TBI of > 6 months duration.** Patients were defined as ambulatory.

### What is moderate to high intensity gait training?

**Moderate to high intensity gait** (or walking) training is **large amounts of** structured and graded walking practice delivered at **moderate to high cardiovascular intensities** with targeted heart rate ranges of 60-80% of heart rate reserve or 70-85% of maximum heart rate.<sup>1</sup>

### Will I harm my patient when delivering high intensity gait training?

**Evidence:** The potential for adverse events with high intensity gait training has been shown to be no greater than conventional therapy.<sup>2-4</sup>

#### **Recommendations:**

- When in doubt check it out! Gain medical clearance from the referring physician or cardiologist if there are concerns (e.g. multiple comorbidities, decreased exercise tolerance)
- Continuously monitor heart rate, blood pressure and oxygen saturation before, during, after exercise at a minimum
- Follow American College of Sports Medicine (ACSM) and American Heart Association (AHA) [recommendations for safe exercise parameters](#)

### Should I be concerned I am causing orthopedic issues (i.e. pain, joint instability, musculoskeletal trauma) when doing high intensity gait training?

**Evidence:** There is negligible evidence in adults in acute-onset CNS injury that high intensity walking exacerbates pain.<sup>5-7</sup>

#### **Recommendations:**

- Utilize bracing as appropriate (e.g. AFO, taping for ankle inversion, Swedish knee cage)
- Modify upper body support and/or increase body weight support (BWS) if patient is experiencing musculoskeletal discomfort
- Patients should wear comfortable shoes for proper support

### Should I be concerned about my patient's movement quality while performing high intensity gait training?

**Evidence:** Prioritization of movement quality during walking training can reduce the amount and intensity of practice, which are key training parameters to facilitate improvement in locomotor ability. Further, prioritization of movement quality does not appear to actually improve movement quality nor improve functional outcomes when compared to treatment strategies focused on the amount and intensity of practice. Evidence demonstrates improvements in walking outcomes regardless of movement errors during practice and that "perfect practice" does not lead to better outcomes. Additionally, data suggests that movement quality improves following high intensity gait training interventions even when quality of movement is not a priority of training.<sup>8-14</sup> These findings are consistent with motor learning literature that identifies the importance of trial and error practice in skill acquisition, and specifically patient recognition of movement errors and the subsequent altering of motor output to increase walking success.<sup>15-19</sup>

#### **Recommendations:**

- Prioritize intensity and amount of practice
- Allow errors in movement to promote motor learning

# FREQUENTLY ASKED QUESTIONS

## CLINICAL PRACTICE GUIDELINE TO IMPROVE LOCOMOTOR FUNCTION FOLLOWING CHRONIC STROKE, INCOMPLETE SPINAL CORD INJURY AND BRAIN INJURY



### What if my patient needs to work on activities like standing balance and transfers?

**Evidence:** High intensity gait training has been shown to improve non-locomotor tasks like transfers and standing balance, without their explicit practice. Further, larger gains in walking outcomes are observed with prioritization of high intensity gait training as compared to practicing non-locomotor tasks.<sup>4,10,20</sup>

#### Recommendations:

- Prioritize high Intensity gait training
- Monitor the effect of high intensity gait training on balance, transfers and other non-locomotor tasks

### Why are body weight supported treadmill training and robotic-assisted training “should not” recommendations?

**Evidence:** It is important to remember that the recommendations of the CPG are applicable to patient populations who have some capacity to walk. Providing additional support through unweighting or robotics could potentially reduce the intensity of practice, a training parameter that is known to be critical. Body weight support devices or robotics may be indicated to allow for non-ambulatory patients to engage in high intensity gait training, although this population was not included in the CPG.<sup>3,18,21,22</sup>

#### Recommendations:

- Maximize the amount and intensity of work during sessions - don't provide assistance that is not needed!
- Avoid the use of body-weight support and robotics unless necessary for safe participation in high intensity gait training

### What additional equipment could be required to perform high intensity gait training safely?

#### Recommendations:

- [Heart rate monitor](#)
- **Walking space or a treadmill**
- [Safety harness system/devices](#) – may be useful to ensure safety

### What if I do not have enough help to deliver high intensity gait training?

**Evidence:** When patients are challenged with large amounts of intensive walking practice, the potential for imbalance and falling increases. Investing in a fall protection harnessing system can be helpful to optimize safety during sessions when additional support from a second person is not available. Rarely are three people required to deliver high intensity gait training, especially considering the lack of prioritization around movement quality.<sup>15,19,24,25</sup>

#### Recommendations:

- Refer to [“how to” videos](#) under the implementation section for more guidance.

### Should I apply high intensity gait training in inpatient rehabilitation?

**Evidence:** The focus of the CPG was on individuals in the chronic stages following stroke, incomplete spinal cord injury, and traumatic brain injury however there is growing body of evidence that indicates high intensity gait training is feasible and beneficial for the sub-acute stroke population.<sup>4,10,23,24</sup> A CPG is currently underway for this patient population, more to come!

#### Recommendation:

- Refer to the [“Intensity Matters” webpage](#) for evidence in sub-acute populations.

# FREQUENTLY ASKED QUESTIONS

## CLINICAL PRACTICE GUIDELINE TO IMPROVE LOCOMOTOR FUNCTION FOLLOWING CHRONIC STROKE, INCOMPLETE SPINAL CORD INJURY AND BRAIN INJURY



### References

1. Hornby TG, Reisman DS, Ward IG, et al. Clinical Practice Guideline to Improve Locomotor Function Following Chronic Stroke, Incomplete Spinal Cord Injury, and Brain Injury. *Journal of neurologic physical therapy : JNPT*. 2020;44(1):49-100.
2. Pang MY, Charlesworth SA, Lau RW, Chung RC. Using aerobic exercise to improve health outcomes and quality of life in stroke: evidence-based exercise prescription recommendations. *Cerebrovascular diseases (Basel, Switzerland)*. 2013;35(1):7-22.
3. Billinger SA, Arena R, Bernhardt J, et al. Physical activity and exercise recommendations for stroke survivors: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2014;45(8):2532-2553.
4. Hornby TG, Holleran CL, Leddy AL, et al. Feasibility of Focused Stepping Practice During Inpatient Rehabilitation Poststroke and Potential Contributions to Mobility Outcomes. *Neurorehabilitation and neural repair*. 2015;29(10):923-932.
5. Tani Y, Otaka Y, Kudo M, Kurayama T, Kondo K. Prevalence of Genu Recurvatum during Walking and Associated Knee Pain in Chronic Hemiplegic Stroke Patients: A Preliminary Survey. *Journal of stroke and cerebrovascular diseases : the official journal of National Stroke Association*. 2016;25(5):1153-1157.
6. Lundström E, Smits A, Terént A, Borg J. Risk factors for stroke-related pain 1 year after first-ever stroke. *European journal of neurology*. 2009;16(2):188-193.
7. Harrison RA, Field TS. Post stroke pain: identification, assessment, and therapy. *Cerebrovascular diseases (Basel, Switzerland)*. 2015;39(3-4):190-201.
8. Holleran CL, Straube DD, Kinnaird CR, Leddy AL, Hornby TG. Feasibility and potential efficacy of high-intensity stepping training in variable contexts in subacute and chronic stroke. *Neurorehabilitation and neural repair*. 2014;28(7):643-651.
9. Leech KA, Kinnaird CR, Holleran CL, Kahn J, Hornby TG. Effects of Locomotor Exercise Intensity on Gait Performance in Individuals With Incomplete Spinal Cord Injury. *Physical therapy*. 2016;96(12):1919-1929.
10. Hornby TG, Holleran CL, Hennessy PW, et al. Variable Intensive Early Walking Poststroke (VIEWS): A Randomized Controlled Trial. *Neurorehabilitation and neural repair*. 2016;30(5):440-450.
11. Mahtani GB, Kinnaird CR, Connolly M, et al. Altered Sagittal- and Frontal-Plane Kinematics Following High-Intensity Stepping Training Versus Conventional Interventions in Subacute Stroke. *Physical therapy*. 2017;97(3):320-329.
12. Lewek MD, Cruz TH, Moore JL, Roth HR, Dhaher YY, Hornby TG. Allowing intralimb kinematic variability during locomotor training poststroke improves kinematic consistency: a subgroup analysis from a randomized clinical trial. *Phys Ther*. 2009;89(8):829-839.
13. Ardestani MM, Henderson CE, Salehi SH, Mahtani GB, Schmit BD, Hornby TG. Kinematic and Neuromuscular Adaptations in Incomplete Spinal Cord Injury after High- versus Low-Intensity Locomotor Training. *Journal of neurotrauma*. 2019;36(12):2036-2044.
14. Ardestani MM, Hornby TG. Effect of investigator observation on gait parameters in individuals with stroke. *J Biomech*. 2020;100:109602.
15. Dobkin B, Apple D, Barbeau H, et al. Weight-supported treadmill vs over-ground training for walking after acute incomplete SCI. *Neurology*. 2006;66(4):484-493.
16. Yagura H, Hatakenaka M, Miyai I. Does therapeutic facilitation add to locomotor outcome of body weight--supported treadmill training in nonambulatory patients with stroke? A randomized controlled trial. *Archives of physical medicine and rehabilitation*. 2006;87(4):529-535.
17. Piira A, Lannem AM, Sørensen M, et al. Robot-assisted locomotor training did not improve walking function in patients with chronic incomplete spinal cord injury: A randomized clinical trial. *Journal of rehabilitation medicine*. 2019;51(5):385-389.
18. Hornby TG, Campbell DD, Kahn JH, Demott T, Moore JL, Roth HR. Enhanced gait-related improvements after therapist- versus robotic-assisted locomotor training in subjects with chronic stroke: a randomized controlled study. *Stroke*. 2008;39(6):1786-1792.
19. Duncan PW, Sullivan KJ, Behrman AL, et al. Body-weight-supported treadmill rehabilitation after stroke. *The New England journal of medicine*. 2011;364(21):2026-2036.
20. Straube DD, Holleran CL, Kinnaird CR, Leddy AL, Hennessy PW, Hornby TG. Effects of dynamic stepping training on nonlocomotor tasks in individuals poststroke. *Physical therapy*. 2014;94(7):921-933.
21. Billinger SA, Boyne P, Coughenour E, Dunning K, Matlage A. Does aerobic exercise and the FITT principle fit into stroke recovery? *Curr Neurol Neurosci Rep*. 2015;15(2):519.
22. Brazg G, Fahey M, Holleran CL, et al. Effects of Training Intensity on Locomotor Performance in Individuals With Chronic Spinal Cord Injury: A Randomized Crossover Study. *Neurorehabilitation and neural repair*. 2017;31(10-11):944-954.
23. Winstein CJ, Gardner ER, McNeal DR, Barto PS, Nicholson DE. Standing balance training: effect on balance and locomotion in hemiparetic adults. *Archives of physical medicine and rehabilitation*. 1989;70(10):755-762.
24. Barbeau H. Locomotor training in neurorehabilitation: emerging rehabilitation concepts. *Neurorehabilitation and neural repair*. 2003;17(1):3-11.
25. Ada L, Dean CM, Morris ME, Simpson JM, Katrak P. Randomized trial of treadmill walking with body weight support to establish walking in subacute stroke: the MOBILISE trial. *Stroke*. 2010;41(6):1237-1242.