



Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

Summary of July 2017 Topic: Vestibular Rehab in Concussion

Dizziness is a common symptom following concussion, according to one abstract occurrence rates are 67-77%. Imbalance, dizziness, visual and vestibular impairments were possible risk factors for prolonged recovery or worse outcomes following concussion.

The abstracts, including a systematic review, investigated whether vestibular therapy would reduce recovery time, improve outcomes, and is effective in the concussion population.

Overall, the research is limited in quality and breadth but suggest that vestibular therapy can be effective in treating this patient population.

The abstracts highlight that testing for vestibular impairments is warranted in the assessment of individuals with concussion. It is also recommended to treat those specific vestibular impairments. The abstracts also found that additional high-quality research is needed to better determine the efficacy of vestibular intervention in concussion management.

Summary of August 2017 Topic: Relaxation/meditation Strategies for Balance & Vestibular Treatment

Abstracts in August looked at possible relaxation and meditation strategies that could be useful in persons with vestibular disorders. While the research is more limited in these treatment options, the abstracts showed possible interventions that may be beneficial.

The first abstract confirmed that Interdisciplinary treatment improves patient coping, functionality, and satisfaction and decreases overall health care utilization in vestibular patients (1). Breathing techniques were shown to be beneficial in two of the abstracts provided. The results suggest that regulation of the breathing pattern may have an influence on disability related to chronic vestibular disease, while proprioception exercises may improve postural control (2). It was found that diaphragmatic breathing subjects, compared to those in the control group, displayed significantly greater heart rate variability and reported feeling less motion sickness during exposure to the virtual reality experience than those in the control group (5).

Yoga practice can significantly reduce cognitive motor interference, which improves balance, by improving allocation and utilization of attentional resources for both balance control and executive cognitive functioning; thus, resulting in better performance under dual task conditions (3).

Finally, a study suggested mindfulness practice can promote effective heart rate regulation, and thereby promote effective recovery after a stressful event for individuals with headache conditions. Moreover, headache conditions may be associated with dysregulated stress recovery, thus more research is needed on the cardiovascular health and stress resilience of headache sufferers (4).



Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

1. Naber CM, Water-Schmeder O, Bohrer PS, Matonak K, Bernstein AL, Merchant MA. Interdisciplinary treatment for vestibular dysfunction: the effectiveness of mindfulness, cognitive-behavioral techniques, and vestibular rehabilitation. *Otolaryngol Head Neck Surg.* 2011 Jul;145(1):117-24.
2. Jáuregui-Renaud K, Villanueva Padrón LA, Cruz Gómez NS. The effect of vestibular rehabilitation supplemented by training of the breathing rhythm or proprioception exercises, in patients with chronic peripheral vestibular disease. *J Vestib Res.* 2007;17(1):63-72.
3. Subramaniam S, Bhatt T. Effect of Yoga practice on reducing cognitive-motor interference for improving dynamic balance control in healthy adults. *Complement Ther Med.* 2017 Feb;30:30-35.
4. Azam MA, Katz J, Mohabir V, Ritvo P. Individuals with tension and migraine headaches exhibit increased heart rate variability during post-stress mindfulness meditation practice but a decrease during a post-stress control condition - A randomized, controlled experiment. *Int J Psychophysiol.* 2016 Dec;110:66-74.
5. Stromberg SE, Russell ME, Carlson CR. Diaphragmatic breathing and its effectiveness for the management of motion sickness. *Aerosp Med Hum Perform.* 2015 May;86(5):452-7.

Summary of September 2017 Topic: Meniere's Disease

The September topic was Meniere's disease and the selected articles aimed to provide evidence to support clinical presentation of Meniere's disease.

One article assessed whether endolymphatic cavity enlargement could be detected in T2 weighted MRI and therefore used to indicate endolymphatic hydrops and confirm a diagnosis Meniere's Disease. The study found that subjects with Meniere's Disease had a statistically significant increased endolymphatic cavity size compared to controls. Therefore, the current protocols using T2 MRI of the temporal bone can be used to help diagnose Meniere's Disease. (2)

Another article used MRI to measure endolymphatic hydrops in order to differentiate between Meniere's Disease and Vestibular Migraine diagnoses. Meniere's Disease showed significant endolymphatic hydrops via 3D-real-IR MRI but not in subjects with Vestibular Migraine. Therefore, this type of MRI can be helpful to rule in/rule out Meniere's Disease or Vestibular Migraine. This is especially useful because the clinical vestibular presentation of the two diagnoses can be similar. (1)

The final study aimed to examine the role of the vestibular system's influence on postural hemodynamics through examination of patients with Meniere's Disease who experience syncope. The study reinforced the function of vestibular system on circulation regulation via the vestibular sympathetic reflex. (3)

These articles support proper diagnosis and therefore appropriate treatment of patients with Meniere's Disease.

- 1) Sun W, Guo P, Ren T, Wang W. Magnetic resonance imaging of intratympanic gadolinium helps differentiate vestibular migraine from Ménière disease. *Laryngoscope.* 2017 Feb 21.
- 2) Keller JH et al. Detection of endolymphatic hydrops using traditional MR imaging sequences. *Am J Otolaryngol.* 2017 Apr 6. pii: S0196-0709(16)30622- 6.

Summary of Monthly Abstract of the Week Topics From July 2017 to present

3) Pyykkö I, Manchaiah V, Zou J, Levo H, Kentala E. Vestibular syncope: A disorder associated with drop attack in Ménière's disease. *Auris Nasus Larynx*. 2017 May 3. pii: S0385-8146(17)30088-3.

Summary of October 2017 Topic: Pain and the Vestibular System

These past month's abstracts looked at the link between the vestibular system and pain. In the first abstract a specific link between the two was discussed. There are remarkable parallel neurochemical phenotypes for inner ear and trigeminal ganglion cells and these afferent channels appear to converge in shared central pathways for vestibular and nociceptive information processing. These pathways share expression of receptors targeted by anti-migraine drugs (1).

The second abstract assessed the incidence of vestibular dysfunction in patients receiving medication for chronic, noncancer pain or other underlying neurologic disorders and to determined associated follow-up therapeutic and diagnostic recommendations. It was found that patients being treated with medications for chronic, noncancer pain or other underlying neurologic disorders may have a higher-than-average incidence of vestibular dysfunction. Baseline assessment and monitoring of the vestibular apparatus may be indicated for these patients (2). Another study determined whether patients with fibromyalgia, compared to age-matched healthy controls, have differences in dynamic posturography, including sensory, motor, and limits of stability despite having a normal clinical neurological examination (3).

A final study sought to determine if reduced head-on-trunk movement alters VOR suppression and gaze accuracy similar to experiments involving normal subjects and if intentionally increasing head and neck movement affects these dynamics. In patients with chronic neck pain, the internal commands issued for combined eye-head movements have large enough amplitudes to create accurate gaze saccades; however, because of increased neck stiffness and viscosity, the head movements produced are smaller, slower, longer, and more delayed than they should be. VOR suppression is disproportionate to the size of the actual gaze saccades because sensory feedback signals from neck proprioceptors are non-veridical, likely due to prolonged coactivation of cervical muscles. The outcome of these changes in eye-head kinematics is head-on-trunk stability at the expense of gaze accuracy. In the absence of vestibular loss, the practical consequences may be dizziness in the short term and imbalance and falls in the long term (4).

1. Balaban CD. Migraine, vertigo and migrainous vertigo: Links between vestibular and pain mechanisms. *J Vestib Res*. 2011;21(6):315-21. doi: 10.3233/VES-2011-0428.

2. Gilbert JW, Vogt M, Windsor RE, Mick GE, Richardson GB, Storey BB, Herder SL, Ledford S, Abrams DA, Theobald MK, Cunningham D, Kelly L, Herring KV, Maddox ML. Vestibular dysfunction in patients with chronic pain or underlying neurologic disorders. *J Am Osteopath Assoc*. 2014 Mar;114(3):172-8. doi: 10.7556/jaoa.2014.034.

3. Jones KD1, King LA, Mist SD, Bennett RM, Horak FB. Postural control deficits in people with fibromyalgia: a pilot study. *Arthritis Res Ther*. 2011 Aug 2;13(4):R127.

Summary of Monthly Abstract of the Week Topics From July 2017 to present

4. Johnston JL, Daye PM, Thomson GT. Inaccurate Saccades and Enhanced Vestibulo-Ocular Reflex Suppression during Combined Eye-Head Movements in Patients with Chronic Neck Pain: Possible Implications for Cervical Vertigo. *Front Neurol*. 2017 Jan 30;8:23. doi: 10.3389/fneur.2017.00023. eCollection 2017.

Summary of November 2017: Whiplash Associated Disorder (WAD)

The November Topic was Whiplash Associated Disorder (WAD). The five articles ranged in focus including oculomotor control, cervical spine proprioception, dizziness, and balance. Common to all of the articles was that the subjects had a whiplash injury.

One systematic review examined oculomotor control after whiplash injury. Overall the results varied but a main finding was that compensatory eye movements were common, especially in smooth pursuits. Based on this finding, it was proposed that this could negatively affect head and eye coordination. (1)

Four articles looked at the efficacy of physical therapy on reducing symptoms from WAD. The first suggested that neck exercises including a behavioral approach could reduce pain, dizziness and improve balance compared to just performing neck exercises. However, this intervention did not resolve these impairments fully. The authors suggested that the effect of specific exercises for the neck, dizziness and balance should be studied. (2)

An article examined the effect of vestibular physical therapy on neck pain and cervical spine range of motion. Vestibular therapy did not decrease pain intensity nor improve range of motion but it clearly did not worsen either of these. (4) Another article examined the potential benefits of vestibular physical therapy on balance and self-perceived dizziness and balance handicap, using the Dizziness Handicap Inventory. Both of these measures improved after six weeks of vestibular PT. (5)

Finally, an article summarized that known mechanisms for dizziness, imbalance, impaired oculomotor control, cervical spine proprioception exists due to cervical afferent dysfunction following a whiplash injury. This article recommended the examination and targeted treatment of these impairments. (3)

Overall, these articles verify the association between dizziness, imbalance and visual disturbances following a whiplash injury and suggest that targeted assessment and treatment of these impairments is important in the rehabilitation of these patients.

1) Ischebeck BK, de Vries J, Van der Geest JN, Janssen M, Van Wingerden JP, Kleinrensink GJ, Frens MA. Eye movements in patients with Whiplash Associated Disorders: a systematic review. *Man Ther*. 2016 Apr;22:122-30. doi: 10.1016/j.math.2015.10.017.

2) Treleaven J, Peterson G, Ludvigsson ML, Kammerlind AS, Peolsson A. *J Orthop Sports Phys Ther*. Balance, dizziness and proprioception in patients with chronic whiplash associated disorders complaining of dizziness: A prospective randomized study comparing three exercise programs. 2017 Jul;47(7):492-502. doi: 10.2519/jospt.2017.7052.

Summary of Monthly Abstract of the Week Topics From July 2017 to present

3) 3) Treleaven J. Dizziness, Unsteadiness, Visual Disturbances, and Sensorimotor Control in Traumatic Neck Pain. *J Orthop Sports Phys Ther* 2017;47(7):492-502. Epub 16 Jun 2017.

doi:10.2519/jospt.2017.7052

4) Hansson EE, Persson L, Malmström EM. Influence of vestibular rehabilitation on neck pain and cervical range of motion among patients with whiplash-associated disorder: a randomized controlled trial. *J Rehabil Med.* 2013 Sep;45(9):906-10. doi: 10.2340/16501977-1197.

5) 5) Ekvall Hansson E, Månsson NO, Ringsberg KA, Håkansson A. Dizziness among patients with whiplash-associated disorder: a randomized controlled trial. *J Rehabil Med.* 2006 Nov;38(6):387-90.

Summary of December 2017: Motion Sensitivity

Motion sickness is a common disturbance occurring in healthy people as a physiological response to exposure to motion stimuli that are unexpected on the basis of previous experience. The motion can be either real, and therefore perceived by the vestibular system, or illusory, as in the case of visual illusion. A multitude of studies has been performed in the last decades, substantiating different nauseogenic stimuli, studying their specific characteristics, proposing unifying theories, and testing possible countermeasures (1).

One study sought to look at the relationship between migraine and motion sickness. Migraine is associated with enhanced motion sickness susceptibility and can cause episodic vertigo, but the mechanisms relating migraine to these vestibular symptoms remain unclear. In this study a hypothesis was tested that the central integration of rotational cues (from the semicircular canals) and gravitational cues (from the otolith organs) is abnormal in migraine patients. They found eye movement responses in Vestibular Migraine patients differed from migraine and normal subjects in three ways: the VOR axis shifts were larger in Vestibular Migraine patients, the normalized axis shift and normalized dumping efficacy were not correlated in VM patients, and the residual conflict in Vestibular Migraine patients was positively correlated with motion sickness susceptibility (2).

Another study compared the rates of carsickness in patients with Vestibular Migraine, Non-vestibular migraine, and Meniere's disease. Overall, 78.4% of the Vestibular Migraine patients had experienced carsickness in their lifetime. 89.2% of the 'definite Vestibular Migraine' patients, and 70.5% of the 'probably Vestibular Migraine' patients had a history of carsickness compared to 43.6% of the Non-Vestibular Migraine patients, and 18.2% of the Meniere's patients. Among the patients who had experienced carsickness, most had experienced carsickness in childhood before the onset of Vestibular Migraine, Non-Vestibular Migraine, or Meniere's disease (3).

For more information the first abstract has a full text with many other studies cited for further review: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4753518/>

1. Bertolini G, Straumann D. Moving in a Moving World: A Review on Vestibular Motion Sickness. *Front Neurol.* 2016 Feb 15;7:14. doi: 10.3389/fneur.2016.00014. eCollection 2016.

Summary of Monthly Abstract of the Week Topics From July 2017 to present

2. Wang J, Lewis RF. Contribution of intravestibular sensory conflict to motion sickness and dizziness in migraine disorders. *J Neurophysiol.* 2016 Oct 1;116(4):1586-1591. doi: 10.1152/jn.00345.2016. Epub 2016 Jul 6.
3. Chang TP, Hsu YC. Vestibular migraine has higher correlation with carsickness than non-vestibular migraine and Meniere's disease. *Acta Neurol Taiwan.* 2014 Mar;23(1):4-10.

Summary of January 2018: Psychological Disorders and Vestibular Dysfunction

January's five abstracts were on the topic of psychological disorders and vestibular dysfunction. One article looked at the correlation of patients with vestibular dysfunction and psychological disorders. This article found that about 20% of this patient population had significant psychological distress, depression and anxiety. The author's conclusion was that consideration of psychological evaluation is warranted in this population. (3)

Another article concluded that vestibular therapy with patients with psychological disorders is warranted and successful. However, these patients may not have as good of outcomes compared to those without psychological disorders. (1) An abstract examined the anxiety and depression severity of patients with acute and chronic dizziness. This article found that patients with chronic dizziness had significantly more anxiety and depression with correlated with a higher emotional subscale of the Dizziness Handicap Inventory. Based on their findings they hypothesized that emotional status may prolong recovery from dizziness. (4)

The next abstract looked at the correlation of anxiety and migraine and found that patients with vestibular migraines had increased anxiety compared to migraines without vertigo. (2). Lastly, an article studied patients with mild brain injury and to see if the post-concussive symptoms could be predicted based on psychological factors assessed by standardized testing. This article found that only affective symptoms could be reliably predicted by psychological factors but postconcussive symptoms of vestibular, somatic, anxiety, depression nor cognitive could not be predicted in this way. (5)

These articles highlight the close connection between vestibular disorders and psychological status and should be assessed by clinicians and appropriately addressed for optimal outcomes.

- 1) MacDowell SG, Trommelen R, Bissell A, Knecht L, Naquin C, Karpinski A. The impact of symptoms of anxiety and depression on subjective and objective outcome measures in individuals with vestibular disorders. *J Vestib Res.* 2017 Nov 10. doi: 10.3233/VES-170627. [Epub ahead of print]
- 2) Kutay Ö et al. Vestibular migraine patients are more anxious than migraine patients without vestibular symptoms. *J Neurol.* 2017 Oct;264(Suppl 1):37-41. doi: 10.1007/s00415-017-8439-6. Epub 2017 Mar 9

Summary of Monthly Abstract of the Week Topics From July 2017 to present

3) Kim SK, Kim YB, Park I-S, Hong SJ, Kim H, Hong SM. Clinical Analysis of Dizzy Patients with High Levels of Depression and Anxiety. *Journal of Audiology & Otology*. 2016;20(3):174-178.

doi:10.7874/jao.2016.20.3.174.

4) Roh KJ, Kim MK, Kim JH, Son EJ. Role of Emotional Distress in Prolongation of Dizziness: A Cross-Sectional Study. *J Audiol Otol*. 2017 Dec;22(1):6-12. doi: 10.7874/jao.2017.00290. Epub 2017 Dec 29.

5) Sullivan KA1, Edmed SL, Greenslade JH, White M, Chu K, Lukin B, Lange RT, Lurie JK. Psychological Predictors of Postconcussive Symptoms Following Traumatic Injury. *J Head Trauma Rehabil*. 2017 Oct 27. doi: 10.1097/HTR.0000000000000347. [Epub ahead of print]

Summary of February 2018: Persistent Postural-Perceptual Dizziness (PPPD)

Persistent postural-perceptual dizziness (PPPD) is a newly defined diagnostic syndrome that unifies key features of chronic subjective dizziness, phobic postural vertigo and related disorders. . It describes a common chronic dysfunction of the vestibular system and brain that produces persistent dizziness, non-spinning vertigo and/or unsteadiness. Once recognised, PPPD can be managed with effective communication and tailored treatment strategies, including specialised physical therapy (vestibular rehabilitation), serotonergic medications and cognitive-behavioural therapy (1).

The second abstract discussed the diagnostic criteria for PPPD. PPPD manifests with one or more symptoms of dizziness, unsteadiness, or non-spinning vertigo that are present on most days for three months or more and are exacerbated by upright posture, active or passive movement, and exposure to moving or complex visual stimuli. PPPD may be precipitated by conditions that disrupt balance or cause vertigo, unsteadiness, or dizziness, including peripheral or central vestibular disorders, other medical illnesses, or psychological distress. PPPD may be present alone or co-exist with other conditions (2). Patients with PPPD also show significantly lower mean scores than normal individuals on conditions 2-6 of the Sensory Organization Test and the composite, and lower than patients recovered from an acute vestibular syndrome on conditions 2-3. Recovered patients had significantly lower mean scores than normal individuals on conditions 4-6 and the composite. Patients with PPPD had the greatest likelihood of abnormal sensory analyses (3).

Recent physiologic and neuroimaging data suggest that greater reliance on visual cues for postural control and dysfunction in central visuo-vestibular networks may be important pathophysiologic mechanisms underlying PPPD. Dysfunctions are thought to involve insular regions that encode recognition of the visual effects of motion in the gravitational field. This final abstract tested for altered activity in vestibular and visual cortices during self-motion simulation. For patients with PPPD, difficulties using visual data to discern the effects of gravity on self-motion may adversely affect balance control, particularly for individuals who simultaneously rely too heavily on visual stimuli. In addition, increased activity in the visual cortex, which correlated with severity of dizziness handicap, may be a neural correlate of visual dependence (4).

Summary of Monthly Abstract of the Week Topics From July 2017 to present

1. Popkirov S, Staab JP, Stone J. Persistent postural-perceptual dizziness (PPPD): a common, characteristic and treatable cause of chronic dizziness. *Pract Neurol*. 2018 Feb;18(1):5-13. doi: 10.1136/practneurol-2017-001809. Epub 2017 Dec 5.
2. Staab JP, Eckhardt-Henn A, Horii A, Jacob R, Strupp M, Brandt T, Bronstein A. Diagnostic criteria for persistent postural-perceptual dizziness (PPPD): Consensus document of the committee for the Classification of Vestibular Disorders of the Bárány Society. *J Vestib Res*. 2017;27(4):191-208. doi: 10.3233/VES-170622.
3. Söhsten E, Bittar RS, Staab JP. Posturographic profile of patients with persistent postural-perceptual dizziness on the sensory organization test. *J Vestib Res*. 2016 Jul 2;26(3):319-26.
4. Riccelli R, Passamonti L, Toschi N, et al. Altered Insular and Occipital Responses to Simulated Vertical Self-Motion in Patients with Persistent Postural-Perceptual Dizziness. *Frontiers in Neurology*. 2017;8:529. doi:10.3389/fneur.2017.00529

Summary of March 2018 Topic: CSM Recap

There were four articles highlighting topics presented at CSM 2018. The first article describes a framework for clinical decision making in the acute setting. TiTrATE is an algorithm for assessing dizziness based on timing, triggers, targeted exam and tests. The goal of TiTrATE is to optimize accurate diagnoses. (1)

The next article addressed the question: Should youth who have sustained a concussion always be symptom free prior to returning to sport? The article established occurrence rates of symptoms such as excessive fatigue, headache and anxiety in the youth general population. The goal was to examine if youth who have who have sustained a concussion might have symptoms that occur typically post-concussion even at baseline. This study established that youth in general have do have post-concussion type symptoms at baseline; therefore, it is not necessarily true that concussed youth should be symptom free prior to return to sport. (2)

The next was a systematic review, included 14 articles, looking at the potential benefits of treating posterior canal BPPV with multiple sessions as well as multiple Epley maneuvers in a single session. The rate of success increased to 100% after 5 sessions and success increased from 84% with one repositioning to 92% after 3 were performed in a single session. Overall, repeating the Epley at least twice and multiple sessions achieved the greatest benefit. (3)

The final article looked at the use of Maddox rod test to assess cyclotorsion. The study demonstrated the subjective test may have a high level of assessor and subject error and results could misinform the clinician's clinical decision making. (4)

1. Newman-Toker DE, Edlow JA. TiTrATE: A Novel, Evidence-Based Approach to Diagnosing Acute Dizziness and Vertigo. *Neurol Clin*. 2015 Aug;33(3):577-99, viii. doi: 10.1016/j.ncl.2015.04.011.
2. Hunt AW, Paniccia M, Reed N, Keightley M. Concussion-Like Symptoms in Child and Youth Athletes at Baseline: What Is "Typical"? *J Athl Train*. 2016 Oct;51(10):749-757. doi: 10.4085/1062-6050-51.11.12. Epub 2016 Nov 11.

Summary of Monthly Abstract of the Week Topics From July 2017 to present

3. Reinink H, Wegner I, Stegeman I, Grolman W. Rapid systematic review of repeated application of the epley maneuver for treating posterior BPPV. *Otolaryngol Head Neck Surg.* 2014 Sep;151(3):399-406. doi: 10.1177/0194599814536530. Epub 2014 May 29.
4. Marsh JD, Durkin MW, Hack AE, Markowitz BB, Cheeseman EW. Accuracy of Double Maddox Rod with Induced Hypertropia in Normal Subjects. *Am Orthopt J.* 2014;64:76-80. doi: 10.3368/aoj.64.1.76.

Summary of April 2018: Exercise, Physical Activity and the Vestibular System

The benefits of exercise physical activity are widely known. This past month's abstracts sought to look at how physical activity may affect the vestibular system.

The first abstract looked at the effect of exercise in migraines. The aim of the study was to investigate the efficacy and possible anti-inflammatory benefits of exercise in patients with vestibular migraine. They found the group undergoing exercise training showed significant symptomatic improvement and demonstrated suppressed antioxidant enzyme activity (1).

The second abstract compared the effects of galvanic vestibular stimulation (GVS) on postural control for participants of different physical activity status - active and non-active. The main results indicated that the regular practice of sports activities counteracts postural control disruption caused by GVS. The active group demonstrated better postural control than the non-active group when subjected to higher vestibular manipulation. The study concluded active participants could identify the relevant sensory input, thought a better central integration, which enables them to switch faster between sensory inputs (2).

Another abstract investigated the relationships between motion sickness susceptibility (MSS) in adulthood and physical and sporting activities (PSA). Subjects having practiced a sport before the age of 18 have less MSS than the other subjects. By practicing PSA, subjects are less dependent on visual input and use vestibular afferences better (3).

The final abstract study aimed to evaluate physical activity-related benefit on vestibular function. They concluded starting physical activity has immediate beneficial effects on the vestibule, in terms of vestibular stimulation mechanisms these effects soon disappear if this activity is stopped (4).

1. Lee YY, Yang YP, Huang PI, Li WC, Huang MC, Kao CL, Chen YJ, Chen MT. Exercise suppresses COX-2 pro-inflammatory pathway in vestibular migraine. *Brain Res Bull.* 2015 Jul;116:98-105. doi: 10.1016/j.brainresbull.2015.06.005. Epub 2015 Jul 4.
2. Maitre J, Paillard T. Postural Effects of Vestibular Manipulation Depend on the Physical Activity Status. *PLoS One.* 2016 Sep 14;11(9):e0162966. doi: 10.1371/journal.pone.0162966. eCollection 2016.
3. Caillet G, Bosser G, Gauchard GC, Chau N, Benamghar L, Perrin PP. Effect of sporting activity practice on susceptibility to motion sickness. *Brain Res Bull.* 2006 Apr 14;69(3):288-93. Epub 2006 Jan 19.
4. Gauchard GC, Vançon G, Gentine A, Jeandel C, Perrin PP. Physical activity after retirement enhances vestibulo-ocular reflex in elderly humans. *Neurosci Lett.* 2004 Apr 22;360(1-2):17-20.



Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

Summary of May 2018: Topic: Visual Considerations in Vestibular Rehabilitation

The May topic was visual considerations in vestibular rehabilitation. The first article developed the Pediatric Visually Induced Dizziness Questionnaire (PVID) to measure the visually induced dizziness in children. Nearly 300 subjects were included, healthy subjects and those with vestibular or migraine disorders. The PVID was found to be a valid measure to identify visually induced vertigo. (1)

The next article examined the connection between migraine and convergence insufficiency. Four cases were used to highlight this connection in cases where patients with migraines reported difficulty reading and persistent eye strain; convergence insufficiency may have played a role. (2)

Next, the study sought information from the Vestibular/Ocular Motor Screening (VOMS) that could indicate a prolonged recovery post-concussion in youth. Based on a retrospective chart review of 167 patients, all VOMS domains except convergence and accommodation deficits were associated with prolonged recovery. It was proposed that the VOMS could be used to help prognosticate recovery in addition to quantify deficits. (3)

The next article aimed to use visual examination as a biomarker for the presence of concussion. Patients with concussion were compared to healthy controls. Average constriction velocity (ACV), average dilation velocity (ADV) and near point convergence were found to significantly affected in those with acute concussion and therefore could be used to help diagnose the presence of concussion. (4)

The final article looked at the role of contrast lighting in the adaptation of the VOR. They found that adaptation was optimized when using a contrast threshold of ~ 1000 during VOR retraining. They note that this optimal level of contrast is far greater than what typical room light provides. (5)

1. Pavlou M, Whitney SL, Alkathiry AA, Huett M, Luxon LM, Raglan E, Godfrey EL, Bamiou DE. Visually Induced Dizziness in Children and Validation of the Pediatric Visually Induced Dizziness Questionnaire. *Front Neurol.* 2017 Dec 5;8:656. doi: 10.3389/fneur.2017.00656. eCollection 2017.
2. Singman EL, Matta NS, Silbert DI. Convergence insufficiency associated with migraine: a case series. *Am Orthopt J.* 2014;64:112-6. doi: 10.3368/aoj.64.1.112.
3. Anzalone AJ, Blueitt D, Case T, McGuffin T, Pollard K, Garrison JC, Jones MT A Positive Vestibular/Ocular Motor Screening (VOMS) Is Associated With Increased Recovery Time After Sports-Related Concussion in Youth and Adolescent Athletes. *Am J Sports Med.* 2017 Feb;45(2):474-479. doi: 10.1177/0363546516668624. Epub 2016 Oct 28
4. Capó-Aponte JE, Beltran TA, Walsh DV, Cole WR, Dumayas JY. Validation of Visual Objective Biomarkers for Acute Concussion. *Mil Med.* 2018 Mar 1;183(suppl_1):9-17. doi: 10.1093/milmed/usx166.
5. Muntaseer Mahfuz M, Schubert MC, Todd CJ, Figtree WVC, Khan SI, Migliaccio AA. The Effect of Visual Contrast on Human Vestibulo-Ocular Reflex Adaptation. *J Assoc Res Otolaryngol.* 2018 Feb;19(1):113-122. doi: 10.1007/s10162-017-0644-6. Epub 2017 Nov 6.

Summary of Monthly Abstract of the Week Topics From July 2017 to present

Summary of June 2018 Topic: Migraine

This past month's studies focused various symptoms associated with migraine and the effects vestibular rehabilitation can have on those symptoms, as well as other disorders associated with migraine. The first study aimed to compare the effects of vestibular rehabilitation on headache and other outcomes relating to dizziness, and the psychological factors in patients with vestibular migraine, patients with dizziness and tension-type headache, and patients without headache. Vestibular rehabilitation contributed to improvement of headache both in patients with vestibular migraine and patients with dizziness and tension-type headache, in addition to improvement of dizziness and psychological factors (1).

The aim of the second study was to compare anxiety disorders in 3 groups: patients with vestibular migraine (VM), patients with migraine but without vertigo (MO) and healthy controls (HC). The study showed that VM patients are significantly more anxious and agoraphobic than MO patients and HC, displaying higher sensitivity to separation and being more prone to seeking medical reassurance (2).

Recent studies have looked at the response of patients with Mal de Debarquement Syndrome (MdDS) to management with migraine prophylaxis, including lifestyle changes and medical therapy. One study found that management of MdDS as vestibular migraine can improve patients' symptoms and increase the quality of life. In addition, nearly all the patients suffering from MdDS had a personal or family history of migraine headaches or had signs or symptoms suggestive of atypical migraine (3).

Currently, there are no known definitive diagnostic tests that can reliably distinguish between Meniere's Disease and Vestibular Migraines, their differentiation is often difficult. Future studies are needed to help adequately distinguish the diagnosis of both diseases (4).

1. Sugaya N, Arai M, Goto F. Is the Headache in Patients with Vestibular Migraine Attenuated by Vestibular Rehabilitation? *Front Neurol.* 2017 Apr 3;8:124. doi: 10.3389/fneur.2017.00124. eCollection 2017.
2. Kutay Ö, Akdal G, Keskinoglu P, Balci BD, Alkin T. Vestibular migraine patients are more anxious than migraine patients without vestibular symptoms. *J Neurol.* 2017 Oct;264(Suppl 1):37-41. doi: 10.1007/s00415-017-8439-6. Epub 2017 Mar 9.
3. Ghavami Y1, Haidar YM1, Ziai KN1, Moshtaghi O1, Bhatt J1, Lin HW1, Djalilian HR1,2. Management of mal de débarquement syndrome as vestibular migraines. *Laryngoscope.* 2017 Jul;127(7):1670-1675. doi: 10.1002/lary.26299. Epub 2016 Oct 12.
4. Tabet P, Saliba I. Meniere's Disease and Vestibular Migraine: Updates and Review of the Literature. *J Clin Med Res.* 2017 Sep;9(9):733-744. doi: 10.14740/jocmr3126w. Epub 2017 Jul 27.

Summary of Monthly Abstract of the Week Topics From July 2017 to present

Summary of July 2018 Topic: Unilateral Vestibular Hypofunction

The topic for July was unilateral vestibular hypofunction. The first article sought to assess the efficacy of vestibular rehabilitation with this patient population. The authors performed a systematic review including 39 articles representing over 2000 patients with unilateral vestibular hypofunction. The evidence was compelling that vestibular rehabilitation is effective in managing patients with unilateral vestibular hypofunction. (1)

The next article looked at the testing to diagnose hypofunction. It compared caloric testing, the historical gold standard, to video head impulse test (vHIT). 324 patients with unilateral hypofunction were given caloric and vHIT. 12% had an abnormal vHIT while 35% had an abnormal caloric testing. A positive vHIT had a high positive predictive value of an abnormal caloric test. Therefore, the vHIT does not replace caloric testing but is a good clinical test for screening of unilateral hypofunction. (2)

The next article aimed to gain information about the functional differences between those with unilateral or bilateral hypofunction and those who did not. 24 subjects were tracked in their gait patterns and eye gaze during stair climbing and descending. Patients with hypofunction looked more at the structure, ie- stairs or ramp. (3)

The final study examined gait mechanic differences between those with unilateral vestibular hypofunction and healthy subjects. Subjects walked on a treadmill with over 30 joint position sensors. The hypofunction group had a wider step width and decreased stability. (4)

1. Is vestibular rehabilitation effective in improving dizziness and function after unilateral peripheral vestibular hypofunction? An abridged version of a Cochrane Review. Hillier S1, McDonnell M. Eur J Phys Rehabil Med. 2016 Aug;52(4):541-56. Epub 2016 Jul 12.
2. Determining vestibular hypofunction: start with the video-head impulse test. van Esch BF1, Nobel-Hoff GE2, van Benthem PP3, van der Zaag-Loonen HJ4, Bruintjies TD4. Eur Arch Otorhinolaryngol. 2016 Nov;273(11):3733-3739. Epub 2016 Apr 25.
3. Patients with chronic peripheral vestibular hypofunction compared to healthy subjects exhibit differences in gaze and gait behaviour when walking on stairs and ramps. Swanenburg J1,2, Bähler E1,3, Adelsberger R4, Straumann D5, de Bruin ED3. PLoS One. 2017 Dec 18;12(12):e0189037. doi: 10.1371/journal.pone.0189037. eCollection 2017.
4. Characterizing Patients with Unilateral Vestibular Hypofunction Using Kinematic Variability and Local Dynamic Stability during Treadmill Walking. Liu P1,2, Huang Q2, Ou Y2, Chen L2, Song R1, Zheng Y2. Behav Neurol. 2017;2017:4820428. doi: 10.1155/2017/4820428. Epub 2017 Jul 13.

Summary of August 2018 Topic: Optometric Examination

This past month's abstracts covered topics related to eye movements that maybe helpful in determining vestibular pathology during examination.

Skew deviation can be defined as vertical misalignment of the eyes that does not map to any of cyclovertical muscles, in association with neurologic symptoms and signs and with posterior fossa lesion.

Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

It is commonly caused by ischemia of the posterior paramedian pons, medial thalamus, or cerebellum. When the vestibular nuclei are involved, skew deviation may occur with acute vestibular syndrome. (1).

In addition to a skew deviation, ocular tilt reaction (OTR) may be present and involves the triad of ocular torsion, skew deviation, and head tilt. Ipsiversive OTR components associated with hearing loss can be early diagnostic signs of anterior inferior cerebellar artery infarction (2)

A review of literature published from January 2016 to August 2017, looked at nystagmus in clinical practice. A structured description of nystagmus including its three-dimensional beating direction, trigger factors, and duration is of major importance. The differential diagnosis of downbeat nystagmus is broad and includes acute intoxications, neurodegenerative disorders and cerebrovascular causes amongst others. In patients with positional nystagmus, the distinction between frequent benign peripheral and rare but dangerous central causes is imperative (3).

Finally, the use of video head impulse test (vHIT) gains and corrective saccades (CSs) at the acute and follow-up stages of vestibular neuritis were assessed. The abnormal rates based on both vHIT gains and CS measurements are similar at the acute stage of VN but are considerably higher at the follow-up stage compared with the abnormal rates based on vHIT gains alone. It is thus advisable to check both CS and vHIT gain while performing vHIT to detect vestibular hypofunction (4)

1. Hernowo A, Eggenberger E. Skew deviation: clinical updates for ophthalmologists. *Curr Opin Ophthalmol*. 2014 Nov;25(6):485-7. doi: 10.1097/ICU.000000000000105.
2. Naoi T, Morita M, Kawakami T, Fujimoto S. Ipsiversive Ocular Torsion, Skew Deviation, and Hearing Loss as Initial Signs of Anterior Inferior Cerebellar Artery Infarction. *Intern Med*. 2018 Jul 1;57(13):1925-1927. doi: 10.2169/internalmedicine.0283-17. Epub 2018 Feb 9.
3. Tarnutzer AA, Straumann D1. Nystagmus. *Curr Opin Neurol*. 2018 Feb;31(1):74-80. doi: 10.1097/WCO.0000000000000517.
4. Yang CJ, Cha EH, Park JW, Kang BC, Yoo MH, Kang WS, Ahn JH, Chung JW, Park HJ. Diagnostic Value of Gains and Corrective Saccades in Video Head Impulse Test in Vestibular Neuritis. *Otolaryngol Head Neck Surg*. 2018 Aug;159(2):347-353. doi: 10.1177/0194599818768218. Epub 2018 Apr 10.

Summary of September 2018 Topic: Vestibular Paroxysmia

The September abstracts' topic was vestibular paroxysmia (VP). One article outlines the diagnostic criteria for VP which requires: ten or episodes vertigo lasting less than 1 minute; stereotyped phenomenology; response to carbamazepine/oxcarbazepine; and not accounted for by another diagnosis. Important diagnostic differentials include Meniere's disease, vestibular migraine, BPPV, stroke, fistula and others. The criteria for probable VP was also described. (2)

Another article described the clinical presentation of 3 pediatrics with VP. Nystagmus can be observed during an episode and MRI may reveal compression. VP Should be considered in pediatrics with multiple brief daily episodes of vertigo. (3)



Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

The first article summarized the treatment of VP which primarily is medication, carbamazepine or similar. There are other medication classes available if the above mentioned is not successful. If the symptoms do not respond to medication, a decompression surgery is available to relieve the pressure of the 8th cranial nerve most commonly caused by a loop of the anterior inferior cerebellar artery. Surgery is reserved for intractable cases. (1)

The final article was a study to evaluate the efficacy of oxcarbazepine with a RCT, 18 patients included. The medication group had a significant reduction in episodes and no adverse side effects were revealed. (4)

1. Brandt T, Strupp M, Dieterich M. Vestibular paroxysmia: a treatable neurovascular cross-compression syndrome. *J Neurol*. 2016 Apr;263 Suppl 1:S90-6. doi: 10.1007/s00415-015-7973-3. Epub 2016 Apr 15.
2. Strupp M, Lopez-Escamez JA, Kim JS, Straumann D, Jen JC, Carey J, Bisdorff A, Brandt T. Vestibular paroxysmia: Diagnostic criteria. *J Vestib Res*. 2016;26(5-6):409-415. doi: 10.3233/VES-160589.
3. Lehnen N, Langhagen T, Heinen F, Huppert D, Brandt T, Jahn K. Vestibular paroxysmia in children: a treatable cause of short vertigo attacks. *Dev Med Child Neurol*. 2015 Apr;57(4):393-6. doi: 10.1111/dmcn.12563. Epub 2014 Aug 22.
4. Bayer O, Brémová T, Strupp M, Hüfner K. A randomized double-blind, placebo-controlled, cross-over trial (Vestparoxy) of the treatment of vestibular paroxysmia with oxcarbazepine. *J Neurol*. 2018 Feb;265(2):291-298. doi: 10.1007/s00415-017-8682-x. Epub 2017 Nov 27.

Summary of October 2018 Topic: ICVR Recap

See the Fall Winter Newsletter

Summary of November 2018 Topic: vHIT

The first article was a review of the literature regarding the video head impulse test (vHIT). The review included 27 articles. One finding is that the video HIT is more sensitive than the clinical head impulse test (cHIT). If the vHIT is negative, caloric testing should be performed to rule out a hypofunction. All canals should be tested in an attempt to isolate the weakness.

The next study aimed to compare the results of vHIT and caloric testing in adolescents. 49 subjects were included in the prospective study, average age was 16. All participants underwent bi-thermal caloric testing with air and vHIT on the same day. Calorics were normal in all cases. 84% had abnormal vHIT and approximately 1/3 of these had multiple canal involvement, of these all had the same ear affected. Posterior canal was most commonly affected. vHIT is a good test for adolescents and children and provides useful information.

The main objective of the next study was to assess if a relationship exists between vHIT and self-reported dizziness. Secondary goal included comparing the vHIT to caloric testing. 115 participants filled



Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

out the Dizziness Handicap Inventory (DHI) and had caloric and vHIT testing on the same day. It was determined that the DHI could not be used to predict outcomes of vHIT. The study also showed that vHIT and caloric testing are not redundant but instead complimentary, providing different functional integrity information of the horizontal canal. The vHIT lacks some sensitivity compared to calorics but is easier for the patient to tolerate and takes less time to complete.

Summary of December 2018 Topic: Concussion

December abstracts looked at the role of the vestibular system in concussion, various baseline measurements and their relationship to one another, and also explored the idea of being symptom free before starting a return to play protocol.

The assessment and treatment of sport-related concussion (SRC) often requires a multifaceted approach. Vestibular dysfunction represents an important profile of symptoms and pathology following SRC, with high prevalence and association with prolonged recovery. The most common vestibular disturbances after SRC include benign paroxysmal positional vertigo, vestibulo-ocular reflex impairment, visual motion sensitivity, and balance impairment. When vestibular dysfunction is identified, there is emerging support for applying targeted vestibular rehabilitation to manage this condition. (1)

In the next study there was a focus on the physiological signs of concussion to help narrow the differential diagnosis of PCS in athletes. The physiological effects of exercise on concussion are especially important for athletes. Some athletes with PCS have exercise intolerance that may result from altered control of cerebral blood flow. Systematic evaluation of exercise tolerance combined with a physical examination of the neurologic, visual, cervical, and vestibular systems can in many cases identify one or more treatable postconcussion disorders. (2)

Collegiate varsity athletes participated in study to assess relationships between various concussion baseline assessments including a demographic questionnaire, graded symptom checklist, neurocognitive assessment, and the Sensory Organization Test (SOT). Reaction time and executive function demonstrated significant relationships with SOT balance performance. These cognitive processes may influence athletes' ability to organize and process higher order information and generate appropriate responses to changes in their environment, with respect to balance and injury risk. (3).

After a concussion, guidelines emphasize that an athlete should be asymptomatic before starting a return-to-play protocol. However, many concussion symptoms are nonspecific and may be present in individuals without concussion. Limited evidence exists regarding the presence of "typical" or preinjury (baseline) symptoms in child and youth athletes. Children and youths commonly experienced symptoms at baseline, including fatigue and nervousness. (4)

1. Mucha A, Fedor S, DeMarco D. Vestibular dysfunction and concussion. *Handb Clin Neurol*. 2018;158:135-144. doi: 10.1016/B978-0-444-63954-7.00014-8.

Summary of Monthly Abstract of the Week Topics From July 2017 to present

2. Leddy J, Baker JG, Haider MN, Hinds A, Willer B. A Physiological Approach to Prolonged Recovery From Sport-Related Concussion. *J Athl Train*. 2017 Mar;52(3):299-308. doi: 10.4085/1062-6050-51.11.08.
3. Vander Vegt CB, Register-Mihalik JK, Ford CB, Rodrigo CJ, Guskiewicz KM, Mihalik JP. Baseline Concussion Clinical Measures Are Related to Sensory Organization and Balance. *Med Sci Sports Exerc*. 2018 Sep 19. doi: 10.1249/MSS.0000000000001789. [Epub ahead of print]
4. Hunt AW, Paniccia M, Reed N, Keightley M. Concussion-Like Symptoms in Child and Youth Athletes at Baseline: What Is "Typical"? *J Athl Train*. 2016 Oct;51(10):749-757. doi: 10.4085/1062-6050-51.11.12. Epub 2016 Nov 11.

Summary of January 2019 Topic: Complementary therapies for balance disorders and management of anxiety

The first article looked at the effect of Tai Chi on balance. This study compared balance outcome measures including Berg, TUG, Functional Reach, 50-foot walk test, 5x sit to stand and ABC. Some then participated in Tai Chi while others in the study did not. Those who performed Tai Chi improved on most of the measures and the control group did not. The average age of participants was 80 years and the Tai Chi program was 16 weeks long. The study concludes that Tai Chi can improve mobility, balance and function, however the n was only 16. (1)

Next, a systematic review was performed to evaluate the effect of Tai Chi on risk of falling. Overall Tai Chi was found to decrease risk of falls especially as the frequency of exercise increased. (2)

Lastly, a review of practice of yoga with or without mindfulness practice was examined based on its effect on stress as measured by physiologic parameters such as BP, HR, cortisol levels. The findings were that practice including postures and yoga poses had a regulating effect on sympathetic nervous system. (3)

1. Bubela D, Sacharko L, Chan J, Brady M. Balance and Functional Outcomes for Older Community-Dwelling Adults Who Practice Tai Chi and Those Who Do Not: A Comparative Study. *J Geriatr Phys Ther*. 2017 Nov 9. doi: 10.1519/JPT.0000000000000153.
2. Huang ZG, Feng YH, Li YH, Lv CS. Systematic review and meta-analysis: Tai Chi for preventing falls in older adults. *BMJ Open*. 2017 Feb 6;7(2):e013661. doi: 10.1136/bmjopen-2016-013661
3. Pascoe MC, Thompson DR, Ski CF. Yoga, mindfulness-based stress reduction and stress-related physiological measures: A meta-analysis. *Psychoneuroendocrinology*. 2017 Dec;86:152-168. doi: 10.1016/j.psyneuen.2017.08.008. Epub 2017 Aug 30.

Summary of February 2019 Topic: Pediatric Concussion

Sport-related concussion (SRC) is a major public health problem. Approximately 90% of SRCs in high school athletes are transient; symptoms recover to baseline within 1 week. However, a small percentage of patients remain symptomatic several months after injury, with a condition known as postconcussion syndrome (PCS). Multiple studies this past month discussed risk factors for prolonged recovery. Vision



Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

and vestibular-related deficits are common after concussion and are associated with prolonged recovery times, substantially impacting the quality of life for children. Vestibular rehabilitation in children with concussion is associated with improvement in symptoms as well as visuovestibular performance (1).

The second abstract discussed clinical variables associated with vestibulo-ocular dysfunction (VOD) detected at initial consultation among pediatric patients. Out of 306 subjects, 30.1% of those with acute SRC (65.0% male, mean age 13.9 years) and 43.0% of those with PCS (41.9% male, mean age 15.4 years) met the criteria for VOD at initial consultation. Independent predictors of VOD at initial consultation included female sex, preinjury history of depression, posttraumatic amnesia, and presence of dizziness, blurred vision, or difficulty focusing at the time of injury. Independent predictors of PCS among patients with acute SRC included the presence of VOD at initial consultation, preinjury history of depression, and posttraumatic amnesia at the time of injury.

They also discussed that VOD at initial consultation is associated with prolonged recovery (2).

Another study showed among youth with SRC, risk for development of PCS was higher in those with a personal and/or family history of mood disorders, other psychiatric illness, and migraine (3).

Finally, ADHD was shown to be an antecedent risk factor for SRC and may contribute to a more complicated course of recovery from SRC (4).

1. Storey EP, Wiebe DJ, D'Alonzo BA, Nixon-Cave K, Jackson-Coty J, Goodman AM, Grady MF, Master CL. Vestibular Rehabilitation Is Associated With Visuovestibular Improvement in Pediatric Concussion. *J Neurol Phys Ther.* 2018 Jul;42(3):134-141.
2. Ellis MJ, Cordingley DM, Vis S, Reimer KM, Leiter J, Russell K. Clinical predictors of vestibulo-ocular dysfunction in pediatric sports-related concussion. *J Neurosurg Pediatr.* 2017 Jan;19(1):38-45. doi: 10.3171/2016.7.PEDS16310. Epub 2016 Sep 30.
3. Morgan CD, Zuckerman SL, Lee YM, King L, Beard S, Sills AK, Solomon GS. Predictors of postconcussion syndrome after sports-related concussion in young athletes: a matched case-control study. *J Neurosurg Pediatr.* 2015 Jun;15(6):589-98. doi: 10.3171/2014.10.PEDS14356. Epub 2015 Mar 6.
4. Iaccarino MA, Fitzgerald M, Pulli A, Woodworth KY, Spencer TJ, Zafonte R, Biederman J. Sport concussion and attention deficit hyperactivity disorder in student athletes: A cohort study. *Neurol Clin Pract.* 2018 Oct;8(5):403-411.

Summary of March 2019 Topic: Vestibular Impairments with Thiamine Deficiency

The topic for Abstract of the Week in March 2019 is vestibular impairments with thiamine deficiency. The first article reviewed literature to identify of oculomotor signs presence prior to onset of encephalopathy due to thiamine deficiency. Decreased horizontal vestibular ocular reflex, horizontal nystagmus, and ophthalmoparesis may occur prior to encephalopathy and may respond to medical treatment of the thiamine deficiency. (1)

Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

Next, a case study was done to examine VHIT and manual HIT in a patient with known thiamine deficiency. The horizontal VOR was decreased and improved with thiamine supplementation. (2)

Lastly, a retrospective study of patients with Wernicke's encephalopathy (WE) was done. Peripheral and central vestibular impairments were identified including gaze evoked nystagmus, ataxia and impaired head impulse testing, specifically involving the horizontal semi-circular canals due to medial vestibular nuclei neurons susceptibility to thiamine deficiency, were prevalent. Therefore, VHIT is useful in examining for suspected WE. (3)

1. Kattah JC. The Spectrum of Vestibular and Ocular Motor Abnormalities in Thiamine Deficiency. *Curr Neurol Neurosci Rep*. 2017 May;17(5):40. doi: 10.1007/s11910-017-0747-9.
2. Kattah JC, Guede C, Hassanzadeh B. The medial vestibular nuclei, a vulnerable target in thiamine deficiency. *J Neurol*. 2018 Jan;265(1):213-215. doi: 10.1007/s00415-017-8670-1. Epub 2017 Nov 15.
3. Lee SH, Kim SH, Kim JM, Tarnutzer AA. Vestibular Dysfunction in Wernicke's Encephalopathy: Predominant Impairment of the Horizontal Semicircular Canals. *Front Neurol*. 2018 Mar 12;9:141. doi: 10.3389/fneur.2018.00141. eCollection 2018.

Summary of April 2019 Topic: Biobehavioral Aspects of Vestibular Rehab

Anxiety is strongly associated with vestibular disorders; however, there is a lack of understanding about how physiotherapists respond to people presenting with anxiety within vestibular rehabilitation. This study aimed to explore physiotherapists' current practice in assessing and treating patients with anxiety in vestibular rehabilitation. Ten Physiotherapists in this study requested tailored training and guidance to enhance their ability to manage patients with anxiety more effectively in vestibular rehabilitation (1).

The next abstract evaluated the evidence for psychological treatments for persistent postconcussion symptoms following mild traumatic brain injury. Counselling or cognitive behaviour therapy have the most support but the evidence remains limited. The best results are seen when postconcussion programs use counselling or cognitive behaviour therapy and are targeted for people with an increased risk of persistent symptoms (2).

The association between depression and benign paroxysmal positional vertigo (BPPV) remains debated. Another study aimed to investigate the risk of BPPV in patients with depressive disorders and found that patients with depressive disorders may have an increased risk of developing BPPV, especially those who have hyperthyroidism and systemic lupus erythematosus (3).

Lastly, it is known that dizziness is frequently encountered in medical practice, often takes a chronic course, and can impair the health-related quality of life (HRQoL). However, results on the extent of this impairment of HRQoL are mixed. 203 patients were evaluated for this using the SF-36. Both the physical (PCS-36) and mental (MCS-36) HRQoL are significantly impaired in patients with dizziness. While the impairment in PCS-36 can be explained by clinical symptoms of the dizziness, MCS-36 impairment is largely associated with psychosocial factors. To improve the patient's overall well-being significantly and permanently doctors have to keep in mind both, the clinical symptoms and the psychosocial factors.

Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

Therefore, in addition to the physical examination doctors should integrate a basic psychological examination into the daily routine with dizziness patients (4).

1. Walker A, Kantaris X, Chambers M. Understanding therapeutic approaches to anxiety in vestibular rehabilitation: a qualitative study of specialist physiotherapists in the UK. *Disabil Rehabil.* 2018 Apr;40(7):829-835. doi: 10.1080/09638288.2016.1277393. Epub 2017 Jan 27.
2. Sullivan KA, Kaye SA, Blaine H, Edmed SL, Meares S, Rossa K, Haden C. Psychological approaches for the management of persistent postconcussion symptoms after mild traumatic brain injury: a systematic review. *Disabil Rehabil.* 2019 Feb 11:1-9. doi: 10.1080/09638288.2018.1558292.
3. Hsu CL, Tsai SJ, Shen CC, Lu T, Hung YM, Hu LY. Risk of benign paroxysmal positional vertigo in patients with depressive disorders: a nationwide population-based cohort study. *BMJ Open.* 2019 Mar 30;9(3):e026936. doi: 10.1136/bmjopen-2018-026936.
4. Weidt S, Bruehl AB, Straumann D, Hegemann SC, Krautstrunk G, Rufer M. Health-related quality of life and emotional distress in patients with dizziness: a cross-sectional approach to disentangle their relationship. *BMC Health Serv Res.* 2014 Jul 22;14:317. doi: 10.1186/1472-6963-14-317.

Summary of May 2019 Topic: Vestibular Schwannoma

The topic for May was Vestibular Schwannoma. The first article looked at the functional effect of persons with vestibular schwannoma who were managed conservatively, without surgery etc. Several balance measures showed inferior balance and mobility of the group with vestibular schwannoma. Despite this variance those with schwannoma still had a low fall risk and supports monitoring this population with balance outcome measures. (1)

The next article examined the efficacy of diagnosis of unilateral vestibular schwannoma using both vHIT and caloric testing. 20/30 subjects with diagnosed vestibular schwannoma demonstrated a hypofunction via calorics but only 10/30 had an abnormal gain indicating a unilateral hypofunction using the vHIT. (2)

The next article evaluated the type of schwannoma's effect on vestibular function. The larger cystic tumors had more of a negative impact on gain. (3)

The final article assessed saccades via a PR score, measure of scatter and refixation of saccades, before and after surgical resection of 36 patients with vestibular schwannomas. Patients with worse pre-op dysfunction had faster compensation after surgery. PR score could be used to evaluate recovery post-op. (4)

1. Low Choy NL, Lucey MM, Lewandowski SL, Panizza BJ. Impacts of small vestibular schwannoma on community ambulation, postural, and ocular control. *Laryngoscope.* 2017 May;127(5):1147-1152. doi: 10.1002/lary.26105. Epub 2016 Aug 13.
2. Tranter-Entwistle I, Dawes P, Darlington CL, Smith PF, Cutfield N. Video head impulse in comparison to caloric testing in unilateral vestibular schwannoma. *Acta Otolaryngol.* 2016 Nov;136(11):1110-1114. Epub 2016 May 25.

Summary of Monthly Abstract of the Week Topics From July 2017 to present

3. Constanzo F, Teixeira BCA, Sens P, Ramina R. Video Head Impulse Test in Vestibular Schwannoma: Relevance of Size and Cystic Component on Vestibular Impairment. *Otol Neurotol*. 2019 Apr;40(4):511-516. doi: 10.1097/MAO.0000000000002158.
4. Batuecas-Caletrio A, Rey-Martinez J, Trinidad-Ruiz G, Matíño-Soler E, Cruz-Ruiz SS, Muñoz-Herrera A, Perez-Fernandez N. Vestibulo-Ocular Reflex Stabilization after Vestibular Schwannoma Surgery: A Story Told by Saccades. *Front Neurol*. 2017 Jan 25;8:15. doi: 10.3389/fneur.2017.00015. eCollection 2017.

Summary of June 2019 Topic: Visual Dependency in Vestibular Disorders

Visual dependence in postural control, often measured by increased postural sway on exposure to visual motion, is an indication of altered visual-vestibular integration with greater weighting towards visual cues for balance. In the first study the relationship between visual dependence in postural control was investigated in relation to cortisol reactivity to psychosocial stress (using the Trier Social Stress Test for groups: TSST-G), as an index of hypothalamic-pituitary-adrenal (HPA) axis function, in healthy young females. In those who exhibited a cortisol response, a negative relationship between stress-induced cortisol reactivity and visual dependence in postural control was observed, since those with the largest cortisol response showed less visual motion induced postural sway (1).

A second study found long-term prognosis after vestibular neuritis is not dependent on the magnitude of the peripheral residual damage. Instead, a combination of visuovestibular psychophysical factors (visual dependence), psychological traits and dysfunctional vestibular perception are relevant. Several functional and structural neuroimaging changes develop after vestibular neuritis, which reflect and underlie the aforementioned psychophysiological and psychological features (2).

Visual dependence can often be present in a patient, although little, if any, measurable pathology is present. Although at times it cannot be accurately measured with either standardized testing or pertinent questionnaires, "hypersensitive" patients have a genuine disease and their symptoms are not of psychiatric origin (3).

The degree to which a person relies on visual stimuli for spatial orientation is termed visual dependency (VD). VD is considered a perceptual trait or cognitive style influenced by psychological factors and mediated by central reweighting of the sensory inputs involved in spatial orientation. VD is often measured with the rod-and-disk test, in which participants align a central rod to the subjective visual vertical (SVV). A final study looked to see what effect torsional nystagmic eye movements may have on individual performance. Using caloric ear irrigation, 3D video-oculography, and the rod-and-disk test, we show that caloric torsional nystagmus modulates measures of VD and demonstrate that increases in tilt after irrigation are positively correlated with changes in ocular torsional eye movements. These findings show that measures of VD can be influenced by oculomotor responses induced by caloric stimulation. The findings are of significance for clinical studies, as they indicate that VD, which often increases in

Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

vestibular disorders, is modulated not only by changes in cognitive style but also by eye movements, in particular nystagmus (4).

1. Smyth N, Flynn M, Rajcani J, F Hucklebridge M, Thorn L, Wood C, Golding J, Evans P, Clow A. Attenuated cortisol reactivity to psychosocial stress is associated with greater visual dependency in postural control. *Psychoneuroendocrinology*. 2019 Jun;104:185-190. doi: 10.1016/j.psyneuen.2019.02.028. Epub 2019 Mar 2.
2. Bronstein AM, Dieterich M. Long-term clinical outcome in vestibular neuritis. *Curr Opin Neurol*. 2019 Feb;32(1):174-180. doi: 10.1097/WCO.0000000000000652.
3. Maire R, Mallinson A, Ceyte H, Caudron S, Van Nechel C, Bisdorff A, Magnusson M, Petersen H, Kingma H, Perrin P. Discussion about Visual Dependence in Balance Control: European Society for Clinical Evaluation of Balance Disorders. *J Int Adv Otol*. 2017 Dec;13(3):404-406. doi: 10.5152/iao.2017.4344.
4. Roberts RE, Da Silva Melo M, Siddiqui AA, Arshad Q, Patel M. Vestibular and oculomotor influences on visual dependency. *J Neurophysiol*. 2016 Sep 1;116(3):1480-7. doi: 10.1152/jn.00895.2015. Epub 2016 Jun 29.

Summary of July 2019 Topic: Vestibulotoxicity

The first article summarized the effects of aminoglycoside ototoxicity including symptoms of ataxia, disequilibrium and oscillopsia. This article also presented the risk factors for developing vestibulotoxicity and the possible dosing implications. (1)

This next was a systematic review that examined the prevalence and the types of adverse effects of aminoglycoside (AG) in order to determine which vestibular tests are best for identifying vestibulotoxicity. 27 studies were included in the review, finding vestibulotoxic side effects ranging from 0 to 60%. Authors hypothesize that VHIT and vestibular evoked myogenic potential are likely better tests for early identification of vestibulotoxicity as AG effects hair cells more. They recommend more prospective studies. (2)

The next was a descriptive article examining the effects of AG on 71 patients. The authors suggest that monitoring vestibulotoxicity is essential in this patient population. (3)

The final article was a review that examined the effects of platinum based chemotherapy on the vestibular system. The effect on the horizontal canal as determined by caloric testing was 0-50% across the studies. The conclusion of the review is that platinum based chemotherapy may pose a significant risk for vestibulotoxicity especially cisplatin and with serial exposure and should be further studied and considered as an adverse effect. (4)

1. Rutka J. Aminoglycoside Vestibulotoxicity. *Adv Otorhinolaryngol*. 2019;82:101-110. doi: 10.1159/000490277. Epub 2019 Jan 15.

Summary of Monthly Abstract of the Week Topics From July 2017 to present

2. Van Hecke R, Van Rompaey V, Wuyts FL, Leysens L, Maes L. Systemic Aminoglycosides-Induced Vestibulotoxicity in Humans. *Ear Hear.* 2017 Nov/Dec;38(6):653-662. doi: 10.1097/AUD.0000000000000458.
3. Handelsman JA. Vestibulotoxicity: strategies for clinical diagnosis and rehabilitation. *Int J Audiol.* 2018 Sep;57(sup4):S99-S107. doi: 10.1080/14992027.2018.1468092. Epub 2018 May 9.
4. Prayuenyong P, Taylor JA, Pearson SE, Gomez R, Patel PM, Hall DA, Kasbekar AV, Baguley DM. Vestibulotoxicity Associated With Platinum-Based Chemotherapy in Survivors of Cancer: A Scoping Review. *Front Oncol.* 2018 Sep 25;8:363. doi: 10.3389/fonc.2018.00363. eCollection 2018.

Summary of August 2019 Topic: Motion Sickness

Motion sickness (MS) is evoked by the conflict among somatosensory, visual, and vestibular input. Last month's abstracts explored possible treatment options for this.

The first study explored the different vestibular physiologic response retention patterns after Coriolis acceleration training in student pilots and extend the results for use with Chinese astronauts in the future. All subjects were exposed to five-day continuous or intermittent Coriolis acceleration training. Retention of the vestibulo-autonomic reaction after vestibular training was different for control subjects and student pilots. Uncoupling patterns between post-rotatory nystagmus and the vestibulo-autonomic reaction may be helpful in the design of clinical rehabilitation plans for balance-disorder patients and for exploration of artificial gravity in future space missions (1).

Virtual reality (VR) immersion often provokes subjective discomfort and postural instability, so called VR sickness. Using transcranial direct current stimulation (tDCS), we aimed to investigate whether VR sickness could be relieved by the modulation of cortical excitability in the temporoparietal junction (TPJ). The amelioration of VR sickness was found by anodal tDCS over the right TPJ and might result from relief of the sensory conflict and/or facilitation of vestibular function (2).

A third study investigated the effects of TENS on MS. Motion sickness symptoms, blood pressure (BP), skin temperature, heart rate (HR), and heart rate variability (HRV) were measured. Saliva samples were collected to analyze the level of stress markers. Cognitive function was evaluated with d2 test prior to and after MS provocation. Sympathetic activity increased but parasympathetic activity decreased during MS. TENS was effective in reducing MS symptoms as well as alleviating cognitive impairment (3).

Pitch head-and-trunk movements during constant velocity rotation are a provocative vestibular stimulus that produces vertigo and nausea. When exposed to this stimulus repeatedly, motion sickness symptoms diminish as the subjects habituate. Acetylleucine is a drug that is used to treat acute vestibular vertigo. In this study, we wanted to ascertain whether this drug (a) lessened motion sickness or delayed habituation; (b) accelerated the recovery following habituation; and (c) whether changes in the subjective vertical accompanied habituation.

Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

Acetylcholine neither reduced the nausea associated with this provocative stimulus, nor hastened the acquisition or retention of vestibular habituation of motion sickness and nystagmus but did find. Subjects showed larger error in the subjective visual vertical after habituation, which indicates that spatial orientation is also affected by vestibular training (4).

1. Wang L, Cao Y, Tan C, Zhao Q, He S, Niu D, Tang G, Zou P, Xing L. Uncoupling VOR and vestibuloautonomic retention to Coriolis acceleration training in student pilots and control subjects. *J Vestib Res.* 2017.
2. Takeuchi N, Mori T, Suzukamo Y, Izumi SI. Modulation of Excitability in the Temporoparietal Junction Relieves Virtual Reality Sickness. *Cyberpsychol Behav Soc Netw.* 2018.
3. Chu H1, Li MH, Juan SH, Chiou WY. Effects of transcutaneous electrical nerve stimulation on motion sickness induced by rotary chair: a crossover study. *J Altern Complement Med.* 2012 May;18(5):494-500. doi: 10.1089/acm.2011.0366. Epub 2012 Apr 26.
4. Clément G, Deguine O, Bourg M, Pavy-LeTraon A. Effects of vestibular training on motion sickness, nystagmus, and subjective vertical. *J Vestib Res.* 2007.

Summary of September 2019 Topic: Back to Basics of Vestibular Therapy

The first article looked at a sample of elderly adults with vestibular impairments, n=57. This retrospective study found vestibular therapy delivered 10 sessions, 2x/week, had a significant positive change on balance, function and dizziness as measured by TUG, gait speed and DHI. (1)

The next article divided 33 healthy adults into 3 groups: 1) no training (control), 2) visual feedback weight shift training (WST) coupled with an active horizontal headshake (HS) activity to elicit a vestibular perturbation, 3) visual feedback WST without HS (NoHS). Six sessions were performed 2x/day every other day, (M-W-F). The Sensory Organization Test (SOT) was performed before and after the treatments. Significant changes in the SOT were found with use of visual feedback training suggesting sensory reweighting adaptation had occurred with the training. (2)

The third study examined residual dizziness and imbalance following repositioning treatment for BPPV. This study was experimental, prospective, randomized, controlled study, n=32, to examine the effects of vestibular therapy on this patient's with BPPV. The subjects were randomly assigned to 2 groups, 1) control group (n=15) performing only repositioning 2) experimental group (n=17) performing the maneuvers and vestibular rehabilitation exercises. The subjects were followed for 6 months and the subjects who were treated with therapy beyond repositioning had significantly decreased dizziness with decreased reoccurrence of BPPV. (3)

Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

The final article examined the relationship between benign paroxysmal positional vertigo (BPPV) and anxiety and assess its association with somatic amplification and health anxiety. 120 subjects were included, 60 with BPPV and 60 healthy (non-BPPV). The Beck Anxiety Inventory (BAI), Short Health Anxiety Inventory (SHAI), and Somatosensory Amplification Scale (SSAS) questionnaires were used. Those in the BPPV group had higher anxiety scores than healthy participants, though the levels of anxiety were normal. The authors suggest that monitoring psychological health may help prevent chronic dizziness. (4)

1. Verdecchia DH, Monzón AM, Urbina Jaimes V, Oliveira FR, Paiva L da S, Carvalho TD de. Patient-Reported and Performance Outcomes Significantly Improved in Elderly Patients with Vestibular Impairment following Rehabilitation: A Retrospective Study. *Journal of Aging Research*. August 2018:1-8.
2. Appiah-Kubi KO, Wright WG. Vestibular training promotes adaptation of multisensory integration in postural control. *Gait Posture*. 2019 Sep;73:215-220. doi: 10.1016/j.gaitpost.2019.07.197.
3. Rodrigues DL, Ledesma ALL, Pires de Oliveira CA, Bahmad F Jr. Effect of Vestibular Exercises Associated With Repositioning Maneuvers in Patients With Benign Paroxysmal Positional Vertigo: A Randomized Controlled Clinical Trial *Otol Neurotol*. 2019 Sep;40(8):e824-e829.
4. Özdilek A, Yalınay Dikmen P, Acar E, Ayanoglu Aksoy E, Özdilek A. Determination of Anxiety, Health Anxiety and Somatosensory Amplification Levels in Individuals with Benign Paroxysmal Positional Vertigo *J Int Adv Otol*. 2019 Jul 26. doi: 10.5152/jao.2019.6874.

Summary of October 2019 Topic: Neural Correlates of Dizziness

It is currently unclear how different brain regions/networks process vestibular information and integrate the information into a unified spatial percept related to somatosensory awareness and whether people with recurrent balance complaints have a neural signature as a trait affecting their development of chronic symptoms of vertigo. By using resting state source localized electroencephalography in non-vertiginous state, electrophysiological changes in activity and functional connectivity of 23 patients with balance complaints are analyzed and compared to healthy subjects. The analyses showed increased alpha2 activity within the posterior cingulate cortex and the precuneus/cuneus and reduced beta3 and gamma activity within the pregenual and subgenual anterior cingulate cortex for the subjects with balance complaints. The study showed that these patients have a neural signature or trait that makes them prone to developing chronic balance problems (1).

In the second abstract using event-related functional magnetic resonance imaging (fMRI) study they investigated how the brain of patients with bilateral vestibular failure (BVF) responds to vestibular stimuli. Using whole brain analysis, group comparisons revealed increased brain activity in early visual cortices (V3) and superior temporal gyrus of patients but there was no significant interaction, i.e. stimulus-response function in these regions were still similar in both groups. Brain activity in these regions during (high)GVS increased with higher dizziness-related handicap scores but was not related to the degree of vestibular impairment or disease duration. The data indicate that perceptible GVS-related

Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

cortical responsivity is not diminished but increased in multisensory (visual-vestibular) cortical regions despite bilateral failure of the peripheral vestibular organ (2).

Vestibular patients occasionally report aggravation or triggering of their symptoms by visual stimuli, which is called visually induced dizziness (VID). After using a resting state fMRI scan they found alterations in the visual and vestibular cortical network in VID patients that could underlie the typical VID symptoms such as a worsening of their vestibular symptoms when being exposed to challenging visual stimuli(3).

Finally a group of six studies that strictly met inclusion criteria were analysed to assess cortical-subcortical correlates of acute clinical features related to VN. The present review clearly reveals that sudden UVF may induce a wide variety of cortical and subcortical responses - with changes in different sensory modules - as a result of acute plasticity in the central nervous system (4).

1. Alsalman O, Ost J, Vanspauwen R, Blaivie C, De Ridder D, Vanneste S. The Neural Correlates of Chronic Symptoms of Vertigo Proneness in Humans. PLoS One 2016, Apr 18;11(4):e0152309. doi: 10.1371/journal.pone.0152309. eCollection 2016.
2. Helmchen C, Rother M, Spliethoff P, Sprenger A. Increased brain responsivity to galvanic vestibular stimulation in bilateral vestibular failure. Neuroimage Clin 2019, Jul 19;24:101942. doi: 10.1016/j.nicl.2019.101942
3. Van Ombergen A, Heine L, Jillings S, Roberts RE, Jeurissen B, Van Rompaey V, Mucci V, Vanhecke S, Sijbers J, Vanhevel F, Sunaert S, Bahri MA, Parizel PM, Van de Heyning PH, Laureys S, Wuyts FL. Altered functional brain connectivity in patients with visually induced dizziness. Neuroimage Clin 2017, Feb 28;14:538-545. doi: 10.1016/j.nicl.2017.02.020. eCollection 2017.
4. Micarelli A, Chiaravalloti A, Schillaci O, Ottaviani F, Alessandrini M. Aspects of cerebral plasticity related to clinical features in acute vestibular neuritis: a "starting point" review from neuroimaging studies. Acta Otorhinolaryngol Ital 2016, Apr;36(2):75-84. doi: 10.14639/0392-100X-642. Epub 2016 Apr 29

Summary of November 2019 Topic: Vestibular Hypofunction

The topic for November's abstracts was Vestibular Hypofunction. The first article was intended to examine the optimal timing for performing dynamic visual acuity (DVA) interventions in those with unilateral hypofunction. DVA protocols were provided for 30 min twice per week for 4 weeks. There were 3 groups that followed this protocol but start times were 2 weeks after onset, 3 weeks and after 1 month after onset of hypofunction. The study demonstrated that earlier intervention had better DVA and aVOR recovery and decreased compensatory saccades. (1)



Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

The next abstract examined whether covert saccades help to maintain dynamic visual acuity in those with bilateral vestibular loss. Frequency, latency, consistency of covert saccade initiation, and gain of covert saccades, residual vestibulo-ocular reflex gain were recorded. No correlation was found between residual vestibulo-ocular reflex gain and dynamic visual acuity. Refixation (covert saccades) were found to improve gaze stability in individuals with bilateral hypofunction and support targeted training in these individuals. (2)

The final article was a case report looking at VOR adaptation after 6 months training in an individual with chronic unilateral vestibular hypofunction, 2 years after onset. The subject performed 15 min/day of incremental VOR adaptation (IVA) with an at home device. The VOR adapted substantially, the patient had subjectively less symptoms and participated more in activities. This case study supports further research of the IVA training in chronic unilateral vestibular hypofunction population. (3)

1. Michel L, Laurent T, Alain T. Rehabilitation of dynamic visual acuity in patients with unilateral vestibular hypofunction: earlier is better. *Eur Arch Otorhinolaryngol.* 2019 Oct 21.
2. Hermann R, Pelisson D, Dumas O, Urquizar C, Truy E, Tilikete C. Are Covert Saccade Functionally Relevant in Vestibular Hypofunction? *Cerebellum.* 2018 Jun;17(3):300-307
3. Rinaudo CN, Schubert MC, Cremer PD, Figtree WVC, Todd CJ, Migliaccio AA. Improved Oculomotor Physiology and Behavior After Unilateral Incremental Adaptation Training in a Person With Chronic Vestibular Hypofunction: A Case Report. *Phys Ther.* 2019 Oct 28;99(10)

Summary of December 2019 Topic: The Effect of Vision on VOR

The topic for December Abstracts was the effect of vision on VOR. The first article looked at the effect of target distance on VOR. 5 different target distances were assessed ranging from 10 cm to 150cm in 18 healthy subjects using vHIT. In these subjects, the gain was significantly increased with the closer target distance. Then 10 subjects were tested in the dark and imagined varying distances. These subjects followed a similar pattern. Therefore, closer targets increases gain. (1)

The next article investigated if convergence increases the gain during when irrigated with cold water, intentionally inducing unilateral hyposensitivity. Their experiments showed that convergence (near viewing) increases the VOR gain during irrigation (hyposensitivity) compared to far viewing, which is the same as without irrigation. Therefore, cold irrigation does not stop the peripheral vestibular pathway. (2)

The goal of the final article was to examine the role of visual impairment and correction on VOR gain. There were 3 groups: 1- refractory error corrected with glasses, 2- refractory error corrected with and 3- no visual impairment. vHIT was used to assess VOR gain. There was no significant difference found between groups or within groups (amount of gain correlated to amount of refractory error). Therefore, no correction of refractory error is required for training VOR. (3)

Summary of Monthly Abstract of the Week Topics From July 2017 to present

1. Castro P, Sena Esteves S, Lerchundi F, Buckwell D, Gresty MA, Bronstein AM, Arshad Q. Viewing Target Distance Influences the Vestibulo-Ocular Reflex Gain when Assessed Using the Video Head Impulse Test. *Audiol Neurootol.* 2018;23(5):285-289. Epub 2018 Dec 11.
2. Tamás LT, Lundberg YW, Büki B. Vergence increases the gain of the human angular vestibulo-ocular reflex during peripheral hyposensitivity elicited by cold thermal irrigation. *J Vestib Res.* 2018;27(5-6):265-270. doi: 10.3233/VES-170629.
3. van Dooren TS. The Video Head Impulse Test and the Influence of Daily Use of Spectacles to Correct a Refractive Error. *Front Neurol.* 2018 Mar 7;9:125. doi: 10.3389/fneur.2018.00125. eCollection 2018.

Summary of January 2020 Topic: BPPV Recurrence

The first study examined recurrence rate and timeline of BPPV. There were two treatment groups randomly assigned to either performing Particle Repositioning Maneuver (PRM) or Brandt Daroff (BD) exercise. 80% of the PRM group had a negative Dix Hall Pike test at 1 week after treatment, opposed to only 25% in the BD group. Of the both groups there was ~35% recurrence rate at 48 months but the time to recurrence was significantly longer in the PRM group. (1)

The second article looked at recurrence rate of BPPV after repositioning. The study included 455 subjects with BPPV, the majority had posterior canal. Telephone follow up revealed a ~11% recurrence rate within 3 years with the highest recurrence occurring in the 51-60 year old demographic. The recurrence of posterior canal and horizontal canal was similar. (2)

The 3rd study aimed to determine if previous canal involvement could be used to predict future canal involvement. For example if the patient initially had R posterior canal BPPV, what is the likelihood that their next occurrence would also be in the R posterior canal? This information could then be used to recommend a home exercise program and subsequent occurrences. The study looked at data from 167 patients and found that recurrences could not be predicted, therefore making a recommendation for home repositioning for a recurrence would likely not be effective. (3)

The final study looked at whether a daily canal reposition procedure (CRP) could prevent recurrence of BPPV. 39 subjects with posterior BPPV were included in the study. 17 performed a daily CRP and 22 in the control group did not. The subjects were followed for 2 years and 41% of the all subjects had a recurrence but there was not significant difference between recurrence rate and onset of symptoms between those who performed CRP and those who did not. The authors concluded that a home program will not affect recurrence of posterior canal. (4)

1. Amor-Dorado JC, Barreira-Fernández MP, Aran-Gonzalez I, Casariego-Vales E, Llorca J, González-Gay MA. Particle repositioning maneuver versus Brandt-Daroff exercise for treatment of unilateral idiopathic

Summary of Monthly Abstract of the Week Topics From July 2017 to present

- BPPV of the posterior semicircular canal: a randomized prospective clinical trial with short- and long-term outcome. *Otol Neurotol.* 2012 Oct;33(8):1401-7. doi: 10.1097/MAO.0b013e318268d50a.
2. Wang YQ, Li JR, Zou SZ, Ding YL. Clinical features and recurrence rate on benign paroxysmal positional vertigo. 2019 Dec;33(12):1185-1188. doi: 10.13201/j.issn.1001-1781.2019.12.017.
 3. Kim HJ, Kim JS. The Patterns of Recurrences in Idiopathic Benign Paroxysmal Positional Vertigo and Self-treatment Evaluation. *Front Neurol.* 2017 Dec 15;8:690. doi: 10.3389/fneur.2017.00690. eCollection 2017.
 4. Helminski JO, Janssen I, Hain TC. Daily exercise does not prevent recurrence of benign paroxysmal positional vertigo. *Otol Neurotol.* 2008 Oct;29(7):976-81. doi: 10.1097/MAO.0b013e318184586d.

Summary of February 2020 Topic: Vestibular Migraine

The topic for February AoW was Vestibular Migraine. The first article aimed to look at migraine associated features in vestibular migraine. 279 subjects with known vestibular migraine in Europe were observed. The average age of migraine onset was nearly 22 years old. The headaches were less than 24 hours in nearly 80% of the cases. Common symptoms were (in order of highest frequency to lowest) nausea, phonophobia, photophobia, vomiting and lightheadedness. Approximately ¼ experienced auras. There was a high familial connection, 67%. Compared to patients with migraine, the vestibular migraine group had a shorter duration of headaches. (1)

The second article performed a review of the literature to summarize the clinical features of vestibular migraine. The authors found a variety of vestibular symptoms can occur and often more than one does. These symptoms can be transient or persistent and include vertigo and dizziness which are often accompanied by headache. Other migraine type features include auditory symptoms, photophobia, phonophobia, osmophobia and visual aura. Anxiety is commonly found with vestibular migraine. Abortive treatments include triptans, vestibular suppressants or anti-nausea medications. Other migraine prophylaxis are useful as well such as antiepileptics, beta blockers and anti-depressants. Diagnosis is made by linking the vestibular and migraine features and treatment is multifactorial including migraine prevention and lifestyle modification as well as treating co-morbidities. (2)

The next article used a questionnaire to examine 252 subjects with vestibular migraine. The features of vertigo, accompanying symptoms, familial history, and migraine precursors were the focus of the study. Age of onset of migraine average was 23 but the average age of onset of vertigo was 38. Over ¾ of the subjects reported internal vertigo and ¼ described external vertigo. Approximately half of subjects had vertigo lasting less than 1 hours with 23% of all subjects reporting vertigo lasting less than 5 min. The other half of subjects had symptoms lasting between 1hr and up to 3 days. Only a few subjects had vertigo symptoms lasting longer than 3 days. The next most common symptom was nausea followed by photophobia, phonophobia, vomiting, palpitations, tinnitus, ear fullness and 4% experienced hearing loss. 70% of all subjects had a known family history of migraines and 67% had a known family history of



Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

vertigo. 42% had motion sickness in youth, a migraine pre-cursor indicating that motion sickness may be predictive of migraine later in life. (3)

The final article reviewed vestibular migraine management, specifically the effects of physical therapy (PT) on vestibular symptoms. Six articles were included and they were inconclusive in demonstrating the effects of PT to minimize vestibular symptoms in patients with VM. The authors recommend more studies, specifically blinded, RTCs. (4)

1. Teggi R et al. Clinical Features of Headache in Patients With Diagnosis of Definite Vestibular Migraine: The VM-Phenotypes Projects. *Front Neurol.* 2018 Jun 5;9:395. doi: 10.3389/fneur.2018.00395. eCollection 2018.
2. Beh SC. Vestibular Migraine: How to Sort it Out and What to Do About it. *J Neuroophthalmol.* 2019 Jun;39(2):208-219
3. Teggi R, et al. Clinical Features, Familial History, and Migraine Precursors in Patients With Definite Vestibular Migraine: The VM-Phenotypes Projects. *Headache.* 2018 Apr;58(4):534-544. doi: 10.1111/head.13240. Epub 2017 Dec 4.
4. Alghadir AH, Anwer S. Effects of Vestibular Rehabilitation in the Management of a Vestibular Migraine: A Review. *Front Neurol.* 2018;9:440. Published 2018 Jun 12.

Summary of March 2020 Topic: Telehealth and Vestibular Rehab

Summary of April 2020 Topic: Telehealth and Vestibular Rehab

May 2020 Topic: Bilateral vestibular Summary of Dysfunction and Cochlear Implant

The topic for May was bilateral vestibular dysfunction and cochlear implant. In the first article, 119 patients with bilateral vestibular loss were studied to determine factors related to falling. The cohort was divided into 2 groups, non-fallers and fallers as defined as ≥ 1 fall in past 12 months. Only DHI score >47 and PSQ >27.5 were found to loosely correlate with fall risk. Other data collected were caloric testing, rotatory chair, vHIT, cVEMP, and hearing function but did not predict falls, nor did age, cause of vestibular loss or duration of loss. (1)

The next article aimed to quantify dizziness in 20 subjects with cochlear implant. Overall the testing found low subjective dizziness per the VRBQ and balance was good per ABC and FGA scores but gait

Summary of Monthly Abstract of the Week Topics From July 2017 to present

speed was abnormal as well as gaze instability and specifically poor performance on SOT condition 5. The conclusions of the authors were that patients should all participate in vestibular rehab given vestibular impairments post cochlear implant. (2)

The third article revealed a new pattern of bilateral hypofunction, bilateral posterior canal dysfunction, as it now can be detected by the vHIT. This was a retrospective study which found this dysfunction most common in the elderly, idiopathic and frequently accompanied by sensorineural hearing loss and posterior downbeat nystagmus with the chief complaint of unstable gait.

1. Dobbels B, Lucieer F, Mertens G, Gilles A, Moyaert J, van de Heyning P, Guinand N, Pérez Fornos A, Herssens N, Hallemans A, Vereeck L, Vanderveken O, Van Rompaey V, van de Berg R. Prospective cohort study on the predictors of fall risk in 119 patients with bilateral vestibulopathy. *PLoS One*. 2020 Mar 9;15(3):e0228768. doi: 10.1371/journal.pone.0228768. eCollection 2020.
2. Murray D, Viani L, Garvan J, Murphy A, Vance R, Simoes-Franklin C, Smith J, Meldrum D. Balance, gait and dizziness in adult cochlear implant users: A cross sectional study. *Cochlear Implants Int*. 2020 Jan;21(1):46-52. doi: 10.1080/14670100.2019.1662978. Epub 2019 Sep 18.
3. Lerchundi F, Laffue AH, Olivier M, Gualtieri FJ. Bilateral posterior semicircular canal dysfunction: a new finding with video head impulse test. *J Neurol*. 2020 Apr 28. doi: 10.1007/s00415-020-09793-5.

Summary of June 2020 Topic: Meniere's Disease

This past month, the abstracts focused on updated treatment for Meniere's disease.

The aim of the first abstract was to investigate current treatment practices and self-reported effectiveness in Ménière's disease. Twenty-four different treatments were reported for Ménière's disease, with dietary modifications, diuretics, and betahistine being the most common. Despite many treatment approaches targeting different proposed pathophysiology for Ménière's disease, all treatments are reported as effective by patients. Intratympanic gentamicin provided the greatest reductions in vertigo count, functional limitations, and work absenteeism, as well as the fewest respondents reporting post-treatment functional limitations (1).

The second abstract compared the efficacy of intratympanic gentamicin injection on vertigo control, drop attacks, and functional level in Ménière's disease patients with and without a history of migraine. They found intratympanic gentamicin appears equally effective in treating vertigo and drop attacks in Ménière's disease with and without migraine, but patients with migraine derive significantly less benefit in terms of functional improvement (2).

The next abstract evaluated the change in quality of life (QOL) of patients with Meniere's disease (MD) after treatment with migraine prophylaxis therapy. The Majority of MD patients who had all failed diuretic therapy responded positively to medications used for migraine prophylaxis, as indicated by a significant improvement in QOL. This study may further suggest a correlation between the



Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

pathophysiologic basis of disease in MD and vestibular migraine. Patients with MD may be successfully managed with medications intended to treat migraine (3).

In the final abstract the authors discussed that the etiology of both MD and migraine is likely multifactorial, further exploration of the association between the two conditions may illuminate how to best manage them in the future. MD is likely a manifestation of cochleovestibular migraine, which occurs as a result of migraine related changes in both the cochlea and vestibule (4).

1. Ward B, Wettstein V, Golding J, et al. Patient Perceptions of Effectiveness in Treatments for Menière's Disease: a National Survey in Italy. *J Int Adv Otol.* 2019;15(1):112-117. doi:10.5152/iao.2019.5758
2. Liu YF, Renk E, Rauch SD, Xu HX. Efficacy of Intratympanic Gentamicin in Menière's Disease With and Without Migraine. *Otol Neurotol.* 2017;38(7):1005-1009. doi:10.1097/MAO.0000000000001460
3. Ghavami Y, Haidar YM, Moshtaghi O, Lin HW, Djalilian HR. Evaluating Quality of Life in Patients With Meniere's Disease Treated as Migraine. *Ann Otol Rhinol Laryngol.* 2018;127(12):877-887.
4. Sarna B, Abouzari M, Lin HW, Djalilian HR. A hypothetical proposal for association between migraine and Meniere's disease. *Med Hypotheses.* 2020;134:109430.

Summary of July 2020 Topic: Suppression Head Impulse Paradigm (SHIMP)

The topic for the Abstract of the Week for the month of July was Suppression Head Impulse Paradigm (SHIMP), a test that may reveal information about the function of the canal rather than the dysfunction. The first article was a retrospective study including 15 patients that were diagnosed with a unilateral vestibular neuritis. The study included an exam of the vestibulo-saccadic interaction to determine recovery of these patients and deduced this information as useful. (1)

The next study aimed to compare SHIMP and Head Impulse Paradigm (HIMP) testing feasibility in adolescents. 29 subjects healthy subjects were tested. During the HIMP, no saccades were seen, consistent with no abnormal function. SHIMP revealed saccades in all subjects. The conclusion was that these testing methods are feasible. (2)

Another study examined the parameters of the HIMP and SHIMP tests. Subjects with vestibular neuritis were tested and found to have compensatory saccades with the HIMP on the affected sides and no saccades on the unaffected side. SHIMP exam showed saccades on the unaffected side and no or weak saccades on the affected side. The conclusion was that SHIMP can evaluate residual vestibular function or compensation in patients with vestibular neuritis. (3)

Summary of Monthly Abstract of the Week Topics From July 2017 to present

The final study also aimed to compare the results of HIMP and SHIMP in patients with acute vestibular neuritis (AVN). A retrospective study of 21 patients with AVN tests were reviewed. The results revealed a correlation between SHIMP & HIMP with SHIMP gain being lower than HIMP and the difference was larger on the affected side. The peak saccadic velocity (PSV) had significant correlation with canal paresis. The authors concluded suggests that SHIMP could be a useful complimentary test in evaluating vestibular function. (4)

1. Manzari L, Tramontano M. Suppression Head Impulse Paradigm (SHIMP) in evaluating the vestibulo-saccadic interaction in patients with vestibular neuritis. *Eur Arch Otorhinolaryngol.* 2020;10.1007/s00405-020-06085-6.
2. Devantier L, Hoskison E, Ovesen T, Henriksen JM. Suppression head impulse paradigm in healthy adolescents - A novel variant of the head impulse test. *J Vestib Res.* 2018;28(3-4):311-317.
3. Chen FY, Zhang YZ, Wu CQ, et al.[The application value of suppression head impulse paradigm in vestibular neuritis] . 2018;32(18):1374-1377. doi:10.13201/j.issn.1001-1781.2018.18.003
4. Park JS, Lee JY, Nam W, Noh S, Chang SO, Kim MB. Comparing the Suppression Head Impulse Paradigm and the Head Impulse Paradigm in Vestibular Neuritis. *Otol Neurotol.* 2020;41(1):e76-e82.

Summary of August 2020 Topic: Postural Threat Influences in Vestibular Function

Standing balance is significantly influenced by postural threat. While this effect has been well established, the underlying mechanisms of the effect are less understood.

The first abstract investigated how vestibulo-spinal reflexes (VSRs) and vestibulo-ocular reflexes (VORs) measured through vestibular evoked myogenic potentials (VEMPs) and video head impulse test (vHIT) outcomes, respectively, are modulated during standing under conditions of increased postural threat. VEMP amplitudes and horizontal and vertical vHIT gains all increased with standing on high surface height conditions. Changes in VEMP amplitudes as well as horizontal vHIT gains were correlated with changes in electrodermal activity. VEMP amplitude also positively correlated with fear . The findings provide support for a potential central modulation of the vestibular nucleus complex through excitatory inputs from neural centers involved in processing fear, anxiety, arousal, and vigilance (1).

The second abstract examined the effect of threat of perturbation, a different form of postural threat, on coupling (cross-correlation, coherence, and gain) of the vestibulo-muscular relationship. In the "No-Threat" conditions, participants stood quietly on a stable surface. In the "Threat" condition, participants' balance was threatened with unpredictable mediolateral support surface tilts. Surface EMG was recorded from bilateral trunk, hip, and leg muscles. Hip and leg muscles exhibited significant increases in peak cross-correlation amplitudes, coherence, and gain in the Threat condition compared with No-Threat conditions, and significant correlations were observed between threat-related changes in physiological arousal and medium-latency peak cross-correlation amplitude in medial gastrocnemius (2).

Summary of Monthly Abstract of the Week Topics From July 2017 to present

The third study repeatedly exposed individuals to height-induced postural threat to determine if reducing the emotional response to threat influences standing balance control. When initially threatened, individuals leaned backward and demonstrated smaller amplitude and higher frequency of COP adjustments; these balance outcomes did not change following repeated threat exposure. Only high frequency COP oscillations (>1.8 Hz) and ankle muscle co-contraction showed any adaptation; regression analyses showed that these behavioural adaptations were accounted for by a combination of emotional and cognitive state changes. This suggests that some threat-induced standing balance changes are more closely linked with the emotional response to threat than others, and are therefore amendable to intervention (3).

Postural threat, known to increase balance-related fear and anxiety, influences vestibular gains during quiet standing in young healthy adults. The final abstract examined whether there is a similar relationship for peripheral unilateral vestibular loss (UVL) patients in comparison to age-matched healthy controls (HC).

By testing subjects center of foot pressure and 90% ranges for angle amplitude and velocity for trunk sway, they provide evidence for a differential effect of height induced postural threat on balance control between UVLs and HCs presumably due to the reduced vestibular-spinal gain in UVL subjects (4).

1. Naranjo EN, Cleworth TW, Allum JHJ, Inglis JT, Lea J, Westerberg BD, Carpenter MG. Vestibulo-spinal and vestibulo-ocular reflexes are modulated when standing with increased postural threat. *J Neurophysiol.* 2016 Feb 1;115(2):833-42. doi: 10.1152/jn.00626.2015. Epub 2015 Dec 2.
2. Lim SB, Cleworth TW, Horslen BC, Blouin JS, Inglis JT, Carpenter MG. Postural threat influences vestibular-evoked muscular responses. *J Neurophysiol.* 2017 Feb 1;117(2):604-611. doi: 10.1152/jn.00712.2016. Epub 2016 Nov 9.
3. Zaback M, Adkin AL, Carpenter MG. Adaptation of emotional state and standing balance parameters following repeated exposure to height-induced postural threat. *Sci Rep.* 2019 Aug 28;9(1):12449. doi: 10.1038/s41598-019-48722-z.
4. Cleworth TW, Allum JHJ, Luu MJ, Lea J, Westerberg BW, Carpenter MG. The Effect of Unilateral Vestibular Loss on Standing Balance During Postural Threat. *Otol Neurotol.* 2020 Aug;41(7):e945-e951. doi: 10.1097/MAO.0000000000002485.

Summary of September 2020 Topic: COVID-19 Neurologic Implications

The topic for Abstract of the Week for September was Neurologic Implications of COVID-19. One article sought to review the literature on clinical otorhinolaryngological presentations due to COVID-19. The main symptoms found were cough, sore throat, shortness of breath. Other symptoms included runny nose, congestion, dizziness and decreased sense of smell. In younger population upper respiratory symptoms were common especially initially when COVID testing could still be negative. Therefore, otolaryngologists should wear proper PPE when examining patients. (1)



Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

The next two articles also reviewed the literature for neurologic symptoms associated with COVID-19. Neurologic symptoms include: headache, dizziness, convulsions and cerebrovascular disease. (3) The systematic review included 61 articles and included 51 articles in the meta-analysis. Smell impairment occurred in 35%, taste impairment in 33%, 19% muscle pain, 12% headache, 10% low back pain, 10% dizziness, 3% cerebrovascular disease and 2% impaired consciousness of patients. Both studies were performed to increase awareness of clinicians treating patients with COVID-19 of these neurologic symptoms and facilitate early diagnosis and treatment. (4)

The final article examined the satisfaction of patients receiving therapy via telehealth across many diagnoses. 205 participants were surveyed about their experiences with overall satisfaction ranging from 93-99% for patient centered outcomes and 86% valued the ability to have telehealth in the future. The pandemic has allowed for widespread use of telehealth which patients value. This is evidence for lawmakers that telehealth should be a benefit extended beyond the pandemic. (2)

1. Krajewska J, Krajewski W, Zub K, Zatoński T. COVID-19 in otolaryngologist practice: a review of current knowledge. *Eur Arch Otorhinolaryngol.* 2020;277(7):1885-1897. doi:10.1007/s00405-020-05968-y
2. Tenforde AS, Borgstrom H, Polich G, et al. Outpatient Physical, Occupational, and Speech Therapy Synchronous Telemedicine: A Survey Study of Patient Satisfaction with Virtual Visits During the COVID-19 Pandemic [published online ahead of print, 2020 Aug 14]. *Am J Phys Med Rehabil.* 2020;10.
3. Gklinos P. Neurological manifestations of COVID-19: a review of what we know so far. *J Neurol.* 2020;267(9):2485-2489. doi:10.1007/s00415-020-09939-5
4. Abdullahi A, Candan SA, Abba MA, et al. Neurological and Musculoskeletal Features of COVID-19: A Systematic Review and Meta-Analysis. *Front Neurol.* 2020;11:687. Published 2020 Jun 26. doi:10.3389/fneur.2020.00687

Summary of October 2020 Topic: Gait

Locomotion involves complex combinations of translational and rotational head movements. The first study investigated horizontal and vertical angular VOR (aVOR) and linear gaze stabilization (IGS) as well as compensation for linear head movements by angular counter rotation of the head during treadmill walking and running at different velocities (0.4 to 2.4 m/s) while fixating either a close (0.5 m) or distant (2.0 m) target. In the horizontal plane, the aVOR predominated throughout all locomotor speeds, whereas the compensation of linear translations was highly variable and generally insufficient. In contrast, in the vertical plane, eye and angular head motion steadily became more in phase with increasing locomotor speed, which served to optimize linear motion compensation. Thus, horizontal and vertical gaze stabilization strategies appear to be considerably different (1).

Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

In lower vertebrates, gaze stabilization during locomotion is at least partially driven by a direct coupling of spinal locomotor commands with extraocular motor signals. The present study thus investigated whether changes in eye-head coordination during human locomotion can be explained by concurrent changes in head movements. Potential correlations between aVOR performance and HMP were analyzed in dependence of locomotor speed and gait cycle phase. They found that the vertical aVOR appears to be suppressed during faster walking and running, whereas at the same time, the predictability of resultant head movements increases. This suggests that during stereotyped human locomotion, internal feed-forward commands supplement or even suppress sensory feedback to mediate gaze stabilization in the vertical plane.

The final study sought to develop a novel sensor-based system to investigate dynamic visual acuity (DVA), walking trajectory, head and trunk movements and the chest-pelvis rotation ratio during forward and backward overground walking in both healthy individuals and patients with vestibular hypofunction. It was demonstrated that individuals with Bilateral Vestibular Hypofunction (BVH) had significant impairments in walking posture and gaze stability using a novel DVA device combined with motion capture. This evaluation may provide detailed and precise information about difficulties patients encounter in their daily lives.

1. Dietrich H, Wuehr M. Strategies for Gaze Stabilization Critically Depend on Locomotor Speed. *Neuroscience*. 2019 Jun 1;408:418-429. doi: 10.1016/j.neuroscience.2019.01.025. Epub 2019 Jan 29.
2. Dietrich H, Wuehr M. Selective suppression of the vestibulo-ocular reflex during human locomotion. *J Neurol*. 2019 Sep;266(Suppl 1):101-107. doi: 10.1007/s00415-019-09352-7. Epub 2019 May 9
3. Chen PY, Chou LW, Jheng YC, et al. Development of a Computerized Device for Evaluating Vestibular Function in Locomotion: A New Evaluation Tool of Vestibular Hypofunction. *Front Neurol*. 2020;11:485. Published 2020 Jun 12. doi:10.3389/fneur.2020.00485

Summary of November 2020 Topic: Current Concepts in Concussion

The topic for Abstract of the week was current concepts in concussion as it relates to vestibular therapy. The first article looked at single and dual task TUG and tandem gait performance in pediatrics with concussion. There were 23 subjects with concussion and 27 controls. All participants performed the protocol twice, average of 10 days later. TUG times were slower in the concussion group in both single and dual task conditions in both first and follow up assessments. Symptoms were significantly worse compared to the control group in the first testing only. Single task tests improved greater than dual task tests and the authors hypothesized that the cognitive component required for dual task execution is delaying improvement. (1)

The next article sought to quantify postural sway in youth with concussion about 7 days after concussion and compared this group to healthy controls. A sensor was used to measure various aspects of sway during single and dual task activities. Significant difference in sway were found between the groups. The

Summary of Monthly Abstract of the Week Topics From July 2017 to present

goal of the study was to determine if postural sway could be used to classify the severity of concussion. Given the significant differences between the groups, the authors concluded that further research is needed to continue to study which aspects of postural sway are most significant to those with concussion. (2)

1. Howell DR, Wilson JC, Brilliant AN, Gardner AJ, Iverson GL, Meehan WP 3rd. Objective clinical tests of dual-task dynamic postural control in youth athletes with concussion. *J Sci Med Sport*. 2019 May;22(5):521-525. doi: 10.1016/j.jsams.2018.11.014. Epub 2018 Nov 22.
2. Bonnette S, Diekfuss JA, Grooms D, Myer GD, Meehan WP 3rd, Howell DR. Integrated linear and nonlinear trunk dynamics identify residual concussion deficits. *Neurosci Lett*. 2020 Jun 11;729:134975. doi: 10.1016/j.neulet.2020.134975. Epub 2020 Apr 13.

Summary of December 2020 Topic: Cognitive Behavior Therapy and Vestibular Therapy for Persistent Dizziness

The December topic for abstract of the week was cognitive behavioral therapy (CBT) and vestibular therapy in treating dizzy patients. The first article proposed that studying combining CBT and vestibular rehab (VR) could be effective in treating dizziness, function and quality of life of people with dizziness. The authors designed a study protocol for future research. The study design proposed was a randomized control trial with 125 subjects from Norway that had been dizzy at least 3 months. The intervention group into treatment group with VR and CBT for 8 sessions or control group with 1 vestibular therapy session. (1)

The next article examined if multiple types of therapy are effective in treating Persistent postural perceptual dizziness (PPPD). The study was retrospective and examined 657 medical records of patients with PPPD and somatic diagnosis. The study showed that patients with PPPD benefit from multiple types of therapies. (2)

The final study examined the long-term efficacy of therapy on treatment of PPPD. 27 patients underwent a treatment program of acceptance and commitment therapy in combination with vestibular therapy 1x/week for 6 weeks. The scores on the Dizziness Handicap Inventory 6 months after treatment. After 6 months, 40% had scores $\leq 14/100$ (remission) and 59% had a reduction in score ≥ 18 . 74% had achieved either remission or reduction in score classified as a treatment effect. Therefore, the authors concluded that combined therapy is possible and may have a long term effect but randomized control studies are needed. (3)

1. Kristiansen, L., Magnussen, L. H., Wilhelmsen, K. T., Mæland, S., Nordahl, S., Clendaniel, R., Hovland, A., & Juul-Kristensen, B. (2019). Efficacy of intergrating vestibular rehabilitation and cognitive behaviour



Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

therapy in persons with persistent dizziness in primary care- a study protocol for a randomised controlled trial. *Trials*, 20(1), 575.

2. Axer, H., Finn, S., Wassermann, A., Guntinas-Lichius, O., Klingner, C. M., & Witte, O. W. (2020). Multimodal treatment of persistent postural-perceptual dizziness. *Brain and behavior*, e01864. Advance online publication.

3. Kuwabara, J., Kondo, M., Kabaya, K., Watanabe, W., Shiraishi, N., Sakai, M., Toshishige, Y., Ino, K., Nakayama, M., Iwasaki, S., & Akechi, T. (2020). Acceptance and commitment therapy combined with vestibular rehabilitation for persistent postural-perceptual dizziness: A pilot study. *American journal of otolaryngology*, 41(6), 102609

Summary of January 2021 Topic: Central Vestibular Disorders

The January topic for abstract of the week was Central Vestibular Dysfunction. The first article looked to examine the structural changes and cortical morphometric features that are associated with migraine and vertigo attacks in patients who have VM. There were 20 patients with VM and 20 normal volunteers who were scanned on an MRI scanner, and their grey matter volume was estimated. The results showed that VM patients have decreased gray matter volume in the prefrontal cortex, posterior insula-operculum regions, inferior parietal gyrus, and supramarginal gyrus. This demonstrates that there are abnormalities in the central vestibular cortex, and there is a linkage between VM and grey matter volume showing that there is a pathophysiological role in these regions in VM patients. (1)

The next article looked at resting state fMRI to look at the changing brain function in patients with persistent postural perceptual dizziness (PPPD). There were 38 patients with PPPD and 38 healthy patients who we compared this data to and also used a machine learning algorithm to classify subjects using the results from the resting state fMRI. The results showed that patients with PPPD showed reduced connectivity in the areas involved in multisensory vestibular processing and spatial cognition, but increased connectivity in areas linking visual and emotional processing. The algorithm showed that dizziness handicap, anxiety, and depression all showed connections in clinically meaningful brain regions. This further shows that connectivity patterns could possibly become a biomarker of PPPD. (2)

The next article looked to examine the treatment of vestibular stimulation in the learning deficits of kids with ASD who are minimally verbal. The researchers conducted a randomized controlled trial with 36 children, 18 of which had ASD and the other 18 were typically developed children. All of the children were trained using a visual-motor Serial-Reaction-Time sequence-learning task in 10 short weekly practice sessions, and a group of children received VS prior to each training session. All of the participants showed reaction time improvement with speed gains, and the VS group showed much larger speed gains compared to those who didn't receive the stimulation and compared to those in the typical development group. The study showed that VS has a positive effect on learning in minimally verbal ASD children. (3)

Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

The last article looked to examine the interaction between the vestibular system and processing and modulating pain. The study looked to use virtual reality to cue the vestibular system using visual cues in interacting with pain. There were 24 healthy volunteers who were placed in a virtual room that appeared to have 5 different degrees of rotation. The results showed that rotating a scenario changed the participants' pain thresholds. This shows that vestibular information that is present in static visual cues can modulate acute pain and that there is research that shows that the visual, vestibular, and somatosensory systems all interact with each other. (4)

1. Zhe, X., Gao, J., Chen, L., Zhang, D., Tang, M., Yan, X., . . . Zhang, X. (2020). Altered structure of the vestibular cortex in patients with vestibular migraine. *Brain and Behavior*, 10(4)
doi:<http://dx.doi.org/10.1002/brb3.1572>
2. Lee, J-O, Lee, E-S, Kim, J-S, et al. Altered brain function in persistent postural perceptual dizziness: A study on resting state functional connectivity. *Hum Brain Mapp.* 2018; 39: 3340– 3353
3. Katz-Nave, G., Adini, Y., Hetzroni, O.E. and Bonne, Y.S. (2020), Sequence Learning in Minimally Verbal Children With ASD and the Beneficial Effect of Vestibular Stimulation. *Autism Research*, 13: 320-337
4. Daniel, A, Barker, L, Martini, M. Pain modulation by illusory body rotation: A new way to disclose the interaction between the vestibular system and pain processing. *Eur J Pain.* 2020; 24: 1119– 1129

Summary of February 2021 Topic: Dizziness and COVID -19

The Abstract of the Week February Topic was dizziness and COVID-19. The first article sought out literature to document the incidence of dizziness with COVID-19. They found 14 articles including 3 case reports and 11 studies of patients citing dizziness or vertigo as a symptom. However, the majority of these articles only noted dizziness as a symptom and did not detail if it was treated or the specific cause of the dizziness ie- labyrinthitis or stroke. The authors emphasize that dizziness is a significant symptom of COVID-19 and should be evaluated by ENT and follow up treatment recommended. (1)

The next article sought to describe the neuro-ophthalmological presentation of coronaviruses. A review of the literature was performed revealing infrequent eye involvement with coronavirus. The most common was viral conjunctivitis but neuro-ophthalmological symptoms included headache, ocular pain, visual disturbances, double vision, cranial nerve palsies, Guillain Barre, encephalitis and nystagmus. The authors recommended that neuro-ophthalmological findings should be evaluated. (2)

The final study sought to describe the neurologic presentation of patients with COVID-19 and other coronaviruses. A systematic review revealed seven studies, a total of 409 patients. The prevalence of neurologic manifestations ranged 17-36%. The most common neurologic finding was headache (16.8%) followed by dizziness (13.9%). The other main neurologic findings were altered consciousness, vomiting, epileptic crisis, neuralgia, hypoxic encephalopathy, acute cerebrovascular disease. (3)

Summary of Monthly Abstract of the Week Topics From July 2017 to present

1. Saniasiaya J, Kulasegarah J. Dizziness and COVID-19. *Ear Nose Throat J.* 2021;100(1):29-30. doi:10.1177/0145561320959573
2. Luís ME, Hipólito-Fernandes D, Mota C, Maleita D, Xavier C, Maio T, Cunha JP, Tavares Ferreira J. A Review of Neuro-Ophthalmological Manifestations of Human Coronavirus Infection. *Eye Brain.* 2020 Oct 30;12:129-137. doi: 10.2147/EB.S268828.
3. Correia AO, Feitosa PWG, Moreira JLS, Nogueira SÁR, Fonseca RB, Nobre MEP. Neurological manifestations of COVID-19 and other coronaviruses: A systematic review. *Neurol Psychiatry Brain Res.* 2020;37:27-32. doi:10.1016/j.npbr.2020.05.008

Summary of March 2021 Topic - General Vestibular Knowledge

The March topic for abstract of the week was General Vestibular Knowledge. The first review looked into the everchanging and range of the topic of vertigo. It examined how previous understanding of “unexplained” dizziness in the elderly is changing, looked into the pathophysiology of vestibular migraine and its relationship to “persistent postural perceptual dizziness”, and discussed how an oculomotor assessment may help identify vestibular presentations of stroke. (1)

The next article looked into vestibular dysfunction and differentiating between peripheral dysfunction vs central dysfunction. To distinguish the two, a full and comprehensive physical examination must be done, and a full history must be obtained from the patient. Once determined if peripheral or central, this will determine the therapeutic approach. The article looked to review different aspects such as history, physical examination, evaluation, epidemiology, differential diagnosis, treatment, complications, and critical points to improve the identification of vestibular dysfunction and differentiating peripheral from central dysfunction. (2)

The next article conducted a literature search examining the current classification and treatment for peripheral, central, and functional vestibular vertigo syndromes. It found that there are internationally accepted diagnostic criteria for BPPV, Menière’s disease, bilateral vestibulopathy, vestibular paroxysmia, and functional dizziness. There are many different treatments for these different pathologies such as: balance is for bilateral vestibulopathy, oxcarbazepine is for vestibular paroxysmia, and vestibular rehabilitation and cognitive behavioral therapy for functional dizziness. This article shows how diagnostically assessing vestibular syndromes has progressed to being much easier now compared to before, but there is still more research that needs to be done when it comes to the treatment of some vestibular syndromes. (3)

The last article looked into the overall role of the vestibular system and how its numerous complex structures and pathways contribute to our sense of proprioception and equilibrium. The central aspect of the vestibular system contains the neural pathways that are responsible for responding to afferent input from the peripheral vestibular system to provide efferent signals to produce our vestibular system



Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

reflexes. There is also data that shows the vestibular system plays a role in consciousness where dysfunctions in the system can cause cognitive deficits related to memory, learning, and navigation. (4)

1. Kaski D. Neurological update: dizziness. *J Neurol*. 2020 Jun;267(6):1864-1869. doi: 10.1007/s00415-020-09748-w. Epub 2020 Mar 4
2. Dougherty JM, Carney M, Emmady PD. Vestibular Dysfunction. 2020 Dec 12. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021
3. Strupp M, Długaiczek J, Ertl-Wagner BB, Rujescu D, Westhofen M, Dieterich M. Vestibular Disorders. *Dtsch Arztebl Int*. 2020 Apr 24;117(17):300-310. doi: 10.3238/arztebl.2020.0300.
4. Casale J, Browne T, Murray I, Gupta G. Physiology, Vestibular System. 2020 May 24. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan. 2020 May 24.

Summary of April 2021 Topic: Exploring Relationship Between Dizziness & Pain

The first article ran 2 studies to examine the relationship between dizziness/vestibular disorders and other medical conditions. The first study sampled 49 patients seeking medical care for balance and dizziness and provided them with a questionnaire about dizziness and pain. 65% of the 49 subjects had both dizziness and pain. The second study sampled seniors (mean age 75) from a fitness center and assessed them for vestibular or balance problems as well as other co-morbidities. 84% of these subjects had a vestibular deficit on testing and 40% also had a cardiovascular co-morbidity, 12% musculoskeletal and 63% reported some other medical problem. Of the 49 subjects, 42% experienced a fall with 1 year of the study start. The authors suggest that examining/addressing vestibular deficits in aging adults is important as well examining for other medical conditions. (1)

The next article further examined the inter-relationship between dizziness and pain. 49 patients referred to a vestibular treatment unit and 62 patients referred to an outpatient psychiatric clinic completed the Dizziness Handicap Inventory. Patients who experienced dizziness had significantly more pain, most commonly located in the head, neck/shoulders and feet. Those with pain also score higher on the emotional component of the DHI. The authors concluded that patients with dizziness or imbalance frequently present with pain which should be examined and treated. (2)

The final study examined the presence of migraine symptoms in patients with Persistent Postural Perceptual Dizziness (PPPD). 19 of 36 total subjects with PPPD were found to have met criteria for migraine headache while 6 met criteria for vestibular migraine. The authors concluded that a majority of patients with PPPD meet criteria for migraine headache which suggests a connection between the two conditions. (3)

Summary of Monthly Abstract of the Week Topics From July 2017 to present

1. Malmström EM, Ekvall Hansson E, Hafström A, Magnusson M, Fransson PA. Co-morbidities to Vestibular Impairments-Some Concomitant Disorders in Young and Older Adults. *Front Neurol.* 2021 Jan 27;11:609928. doi: 10.3389/fneur.2020.609928.
2. Malmström, E. M., Magnusson, M., Holmberg, J., Karlberg, M., & Fransson, P. A. (2020). Dizziness and localized pain are often concurrent in patients with balance or psychological disorders. *Scandinavian journal of pain*, 20(2), 353–362
3. Sarna, B., Risbud, A., Lee, A., Muhonen, E., Abouzari, M., & Djalilian, H. R. (2021). Migraine Features in Patients with Persistent Postural-Perceptual Dizziness. *The Annals of otology, rhinology, and laryngology*, 34894211007233. Advance online publication.

Summary of May 2021 Topic: Technology in Vestibular Rehabilitation

Summary of June 2021 Topic: Peripheral Vestibular Dysfunctions and Functional Outcome Measures

The June topic for abstract of the week was Peripheral Vestibular Dysfunctions and Functional Outcome Measures. The first article looked to examine walking and turning performance during the TUG test in patients with vestibular deafferentation compared to those without and examine to see if the data improved in patients with vestibular deafferentation following vestibular physical therapy. 26 patients with reports of dizziness not from a vestibular origin and 12 healthy controls were recruited and asked to complete the TUG as fast and safely as possible. Results showed that the turning performance in patients with vestibular deafferentation before therapy was significantly more impaired than the healthy controls. After therapy, turning performance was significantly improved and was normalized as compared to the healthy controls. With the positive results, this shows that the TUG can be used to quantify the change in turning performance in patients with vestibular deafferentation after physical therapy. (1)

The next article looked to examine the relationship between the ABC scale and Dizziness Handicap Inventory with balance performance and fall status in patients with peripheral vestibular disorders. This was a retrospective study done in an outpatient balance clinic with 97 patients with dizziness or imbalance symptoms. Results showed that patients with a low-level confidence in their balance and a severe score on the DHI had poorer balance and were more likely to experience multiple falls. The two measures showed strong validity with ABC showing better concurrent validity with balance and DHI better with fall history. (2)

The next article looked at the measures of Functional Gait Assessment and quality of life measurements in patients before and after vestibular schwannoma resection and to assess how initial Functional Gait

Summary of Monthly Abstract of the Week Topics From July 2017 to present

Assessment can predict post-op quality of life. The article took patients older than 18 who underwent the resection where Functional Gait Assessment was administered a week before and after surgery, and the quality of life measure was administered preoperatively and 3 months post-op. Results showed with a statistically significant p-value that pre-op FGA correlated with post-op quality of life measure, but post-op FGA scores did not correlate with post-op quality of life measures. From here, it was determined that although FGA and QOL data showed correlation in the preoperative setting; this was not found postoperatively. More research is needed to determine contributions to post-op quality of life. (3)

The last article looked to examine the different processes that could be used to quantify functional deficits and track improvement in balance following vestibular loss. The 3 processes included: partial peripheral recovery of sensory responses eliciting canal or otolith driven vestibular reflexes, central compensation of vestibular reflex gains, and sensory substitution of visual and proprioceptive inputs for vestibular contributions to balance control. Results showed otolith responses were unaffected, but decreases in all canal driven vestibular ocular reflex gains were observed. Balance control was normal, but VOR gains remained low. This shows how difficult it is to predict balance improvements this way compared to the need to emphasize testing balance control directly to determine true improvements. (4)

1. Kim KJ, Gimmon Y, Millar J, Brewer K, Serrador J, Schubert MC. The Instrumented Timed Up and Go Test Distinguishes Turning Characteristics in Vestibular Hypofunction. *Phys Ther*. 2021 Mar 26:pzab103. doi: 10.1093/ptj/pzab103.
2. Herrensens N, Swinnen E, Dobbels B, Van de Heyning P, Van Rompaey V, Hallems A, Vereeck L. The Relationship Between the Activities-Specific Balance Confidence Scale and Balance Performance, Self-Perceived Handicap, and Fall Status in Patients With Peripheral Dizziness or Imbalance. *Otol Neurotol*. 2021 Apr 14. doi: 10.1097/MAO.0000000000003166. Epub ahead of print.
3. Said M, Lee J, Moshtaghi O, Saliba J, Richardson AJW, Ngo V, Mehranpour P, Schwartz MS, Friedman RA. The Relationship Between the Functional Gait Assessment and Quality-of-Life Data in Patients Undergoing Vestibular Schwannoma Resection. *Otol Neurotol*. 2021 Mar 19. doi: 10.1097/MAO.0000000000003137. Epub ahead of print.
4. Allum JHJ, Rust HM, Honegger F. Functional Testing of Vestibulo-Spinal Contributions to Balance Control: Insights From Tracking Improvement Following Acute Bilateral Peripheral Vestibular Loss. *Front Neurol*. 2019 May 28;10:550. doi: 10.3389/fneur.2019.00550.

Summary of July 2021 Topic: Motor Learning and the Vestibular System

The July topic featured articles co-authored by Dr Michael Schubert, PT, PhD. The first article looked at incremental training of VOR. There were 5 protocols, each performed in 1 single session by 11 patients. Here were the 5 protocols: 1. Unilateral incremental VOR adaptation for 15 min with head impulses equally to each side 2. The head impulse training ratio was increased on the adapting side by 2:1 for 15

Summary of Monthly Abstract of the Week Topics From July 2017 to present

min. 3. All impulses were toward the adapting side for 7.5 min. 4. All impulses toward the adapting side for 15 min. 5. All impulses were toward the adapting size and duration of the visual target exposure was doubled. Active and passive VOR were measured before and after each protocol. The results showed that the amount of training time and visual target exposure affected the amount of gain adaptation achieved suggesting that quality and duration of the performance is more important than number of repetitions. (1)

The next article looked at the role of convergence on the VOR gain adaptation. 22 subjects with unilateral hypofunction performed VOR adaptation exercises at a distance of 15cm and 150cm. Results were compared to 12 health subjects. Distance did not affect the amount of gain adaptation. However, it was found that retinal slip was increased at near distances with rotation toward the side of the lesion as compared to far viewing. In addition, the VOR gain adaptation was increased contralateral to the lesion in the near viewing compared to far viewing. This suggests that adaptation for near viewing may warrant increased ipsilateral exercises since retinal slip is increased with near viewing. (2)

The final article sought to examine the effects of incremental VOR training in subjects with chronic peripheral vestibular hypofunction. This was a double blinded randomized controlled study with 24 subjects with vestibular hypofunction, 13 intervention group and 11 in the control group. The control group performed viewing x1 and the intervention group performed incremental VOR exercises daily for 15 minutes for 4 sessions within a 7 day time frame. VOR was measured with video-oculography. Other measures were compensatory saccades, DVA, static balance, gait and subject symptoms. The intervention group gain increased and the control group did not. The control group had decreased compensatory saccade latency. Both groups had improved DVA. The incremental group also had improvements in some dynamic gait index scores and DHI scores. The authors suggest that incremental VOR training has better training effects compared to x1 viewing. (3)

1. Muntaseer Mahfuz M, Schubert MC, Figtree WVC, Todd CJ, Migliaccio AA. Human Vestibulo-Ocular Reflex Adaptation Training: Time Beats Quantity. *J Assoc Res Otolaryngol*. 2018 Dec;19(6):729-739. doi: 10.1007/s10162-018-00689-w. Epub 2018 Sep 24.
2. Rinaudo CN, Schubert MC, Cremer PD, Figtree WVC, Todd CJ, Migliaccio AA. Once-Daily Incremental Vestibular-Ocular Reflex Adaptation Training in Patients With Chronic Peripheral Vestibular Hypofunction: A 1-Week Randomized Controlled Study. *J Neurol Phys Ther*. 2021 Apr 1;45(2):87-100. doi: 10.1097/NPT.000000000000034
3. Rinaudo CN, Schubert MC, Cremer PD, Figtree WVC, Todd CJ, Migliaccio AA. Once-Daily Incremental Vestibular-Ocular Reflex Adaptation Training in Patients With Chronic Peripheral Vestibular Hypofunction: A 1-Week Randomized Controlled Study. *J Neurol Phys Ther*. 2021 Apr 1;45(2):87-100. doi: 10.1097/NPT.0000000000000348

Summary of Monthly Abstract of the Week Topics From July 2017 to present

Summary of August 2021 Topic: Motor Learning and the Vestibular System

The August topic for abstract of the week was Motor Learning and the Vestibular System. The first article looked to examine how dizzy symptoms interfere with locomotor learning. 20 patients with vestibular neuritis in the chronic phase and 15 patients with Meniere's disease were compared to 15 healthy individuals by measuring locomotor adaptive learning using the "broken escalator" effect. Results showed that the size of the locomotor aftereffect was related to how symptomatic patients were, but the degree of vestibular loss was not related with symptom load or locomotor aftereffect size. From this, it was found that dizziness symptoms influence locomotor adaptation. (1)

The next article looked to examine the influence of dance training on the VOR suppression during passive head impulses. VOR and VOR suppression were assessed in 12 controls and 12 dancers. The results showed that dancers showed a significantly reduced VOR gain in comparison to controls especially with dancers who had been dancing for more than 10 years. This shows that dance training improves VOR suppression and modulates VOR suppression abilities. (2)

The next article looked at the relationship between separate and distinct brain regions and in vivo networks to reveal the metabolic connectome in a rat. Results showed that there were changes in the metabolic brain connectivity. There was an increase in connectivity that occurred within brain regions that were associated with brainstem-cerebellar and vestibular networks. It was shown that quick changes were found in network organization which can show an initial functional substitution of vestibular loss which then show recalibration and reorganization of sensorimotor networks during VC. (3)

The last article looked to examine the brain's changes in response to learning a balance task with their eyes open versus eyes closed. The eyes open group performed better with this task specific test suggesting that brain changes in response to learning a task differ depending on the learning success and availability of visual input and not just independently on the time spent on learning. (4)

1. Patel, M, Roberts, E, Arshad, Q, Bunday, K, Golding, JF, Kaski, D, Bronstein, AM. The "broken escalator" phenomenon: Vestibular dizziness interferes with locomotor adaptation. *J Vestib Res.* 2020; 30(2):81-94. doi: 10.3233/VES-200693.
2. Maheu M, Behtani L, Nooristani M, Delcenserie A, Champoux F. Enhanced vestibulo-ocular reflex suppression in dancers during passive high-velocity head impulses. *Exp Brain Res.* 2019;237(2):411-416. doi:10.1007/s00221-018-5431-z
3. Grosch M, Lindner M, Bartenstein P, et al. Dynamic whole-brain metabolic connectivity during vestibular compensation in the rat. *Neuroimage.* 2021;226:117588. doi:10.1016/j.neuroimage.2020.117588



Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

4. Dordevic M, Taubert M, Müller P, Kaufmann J, Hökelmann A, Müller NG. Brain Gray Matter Volume Is Modulated by Visual Input and Overall Learning Success but Not by Time Spent on Learning a Complex Balancing Task. *J Clin Med*. 2018;8(1):9. Published 2018 Dec 21. doi:10.3390/jcm8010009

Summary of September 2021 Topic: Vestibular Implications in Multiple Sclerosis

October 2021

We apologize for the pause in recent Abstracts. This was due to an error with the group we use to publish these, and it is now corrected. The original topic that was planned for October will now be November's topic. The Abstract below was originally sent the first week of October but will be repeated today. The following weeks will be all new abstracts on concussion. Thank you for your ongoing support of Abstract of the Week!

Summary of November 2021 Topic: Concussion

The November topic for abstract of the week was Concussion. The first article looked to examine the reasoning behind why there is a difference in outcome measures between women and men after a mild TBI. It was seen that women had worse functional outcome measures 6 months after injury and wanted to investigate if this was due to psychiatric history, gender related sociodemographic variables, or by care pathways. There were 1842 men and 1022 women that were included where natural effects models were used to look at the effect of gender on the outcomes. It was shown that the worse outcomes that were seen in women were partly due to psychiatric history and not explained by the other mentioned outcomes. (1)

The next article looked to determine the discriminative validity of Vestibular/ Ocular Motor Screening (VOMS) item scores and overall VOMS score for identifying college athletes in the first few days following a sports-related concussion. 285 athletes who were concussed and 285 healthy controls completed the VOMS within 3 days of injury. It was shown that if the vertical saccades score is greater than or equal to 1 and if the horizontal vestibular/ ocular reflex is greater than or equal to 2 that these were the best discriminates to determine concussion from control. Near point of convergence did not significantly show concussion from control group. This shows that individual VOMS items and overall VOMS scores are useful in revealing concussion in collegiate athletes within 3 days of injury and can be used by clinicians to help identify concussion. (2)

The next article looked to examine the effectiveness of near point of convergence testing following concussion compared to previous measures before concussion and to look at the effectiveness of vision therapy on correcting near point of convergence. There was moderate evidence found that patients do



Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

have impaired near point convergence up to several months post-concussion, and there was low level evidence shown that the impairments can be treated with vision therapy. (3)

The last article looked to examine the relationship between risk factors and vestibular-oculomotor outcomes after sport-related concussion and to see if there was any association between the two. 85 athletes were assessed where they completed a clinical interview, history questionnaire, symptom inventory, and a VOMS screening. It was found that female sex, on-field dizziness, fogginess, and post-traumatic migraine symptoms including headache, nausea, light sensitivity, and noise sensitivity were predictive of experiencing vestibular-oculomotor symptoms or impairment after a sport-related concussion. (4)

1. Mikolic A, Oude Groeniger J, Zeldovich M, Wilson L, Roeters van Lennep JE, van Klaveren D, Polinder S. Explaining outcome differences between men and women following mild traumatic brain injury. *J Neurotrauma*. 2021 Oct 7. doi: 10.1089/neu.2021.0116. Epub ahead of print. PMID: 34617454.
2. Anthony P Kontos 1 2, Shawn R Eagle 1 2, Gregory Marchetti 3 2, . Discriminative Validity of Vestibular Ocular Motor Screening in Identifying Concussion Among Collegiate Athletes: A National Collegiate Athletic Association-Department of Defense Concussion Assessment, Research, and Education Consortium Study. *Am J Sports Med*. 2021 Jul;49(8):2211-2217.
3. Santo AL, Race ML, Teel EF. Near Point of Convergence Deficits and Treatment Following Concussion: A Systematic Review. *J Sport Rehabil*. 2020 Nov 1;29(8):1179-1193. doi: 10.1123/jsr.2019-0428. Epub 2020 Mar 4
4. Womble MN, McAllister-Deitrick J, Marchetti GF, Reynolds E, Collins MW, Elbin RJ, Kontos AP. Risk Factors for Vestibular and Oculomotor Outcomes After Sport-Related Concussion. *Clin J Sport Med*. 2021 Jul 1;31(4):e193-e199. doi: 10.1097/JSM.0000000000000761.

Summary of December 2021 Topic: Postural Threat

Summary of January 2022 Topic: Horizontal Canal BPPV

The January topic for the abstract of the month was Horizontal Canal BPPV. The first article looked to examine the relationship between specific nystagmus parameters in the supine roll test and how they correlate with prognosis after repositioning for horizontal canal BPPV. Patients diagnosed with horizontal canal BPPV were examined through video nystagmography and then used the canalith repositioning procedure for treatment. From here, patients were grouped into the cured group or the resistant group. It was found that of the 813 patients 99 of them were grouped into the resistant group, and these patients showed longer latency in the affected side and slower phase velocity in the unaffected side. From this, it can be concluded that horizontal canal BPPV patients with a longer latency

Summary of Monthly Abstract of the Week Topics From July 2017 to present

in the affected side and slower phase velocity in the unaffected side have a higher risk of BPPV persisting after repositioning. (1)

The next article looked to determine what factors are the most accurate in determining the involved side in horizontal canal BPPV from the bedside. 44 video clips were included observing the nystagmus induced during supine head roll test from patients with horizontal canal BPPV, and the participants were able to watch the videos twice and were to write where the lesion was according to the video. Absolute difference and asymmetry of nystagmus intensity were calculated. After the first watching of the video, the accuracy of bedside was 83.5% and after the second video, the accuracy was 86% with the accuracy being more closely correlated with the asymmetry compared to the absolute difference. This shows that the accuracy of bedside horizontal canal BPPV of the affected side is acceptable when the nystagmus asymmetry is utilized. (2)

The next article looked to examine the effectiveness of the forced prolonged position of liberatory maneuvers in the treatment of lateral canal benign paroxysmal positional vertigo. There were 116 patients assigned to the treatment group of the prolonged position, and there were 105 subjects assigned to the sham treatment. After treatment, these patients were followed up with the supine roll test. Results showed at the follow-up, the effectiveness of the prolonged position compared to the sham maneuver was 57.8% compared to 12.4%. This shows that the forced prolonged position was more effective than the sham maneuver in treating lateral canal benign paroxysmal positional vertigo. (3)

The last article discussed the prevalence of BPPV and the best ways to treat it. BPPV is the most common type of peripheral vertigo that can either present as canalolithiasis or cupulolithiasis. Canalolithiasis happens when the otoconia move freely into one of the semicircular canals, while cupulolithiasis occurs when the otoliths adhere to the cupula. The posterior canal is the most common, while the lateral canal follows behind. The Zuma maneuver and the modified Zuma maneuver have been adopted for the management of lateral canal BPPV. (4)

1. Yu J, Gu Y, Meng G, et al. Nystagmus Parameters of Supine Roll Test Correlates With Prognosis After Repositioning Maneuver in Horizontal Semicircular Canal Benign Paroxysmal Positional Vertigo. *Front Neurol.* 2021;12:790430. Published 2021 Dec 6. doi:10.3389/fneur.2021.790430
2. Choi SY, Oh SW, Kim HJ, Kim JS. Determinants for bedside lateralization of benign paroxysmal positional vertigo involving the horizontal semicircular canal. *J Neurol.* 2020;267(6):1709-1714. doi:10.1007/s00415-020-09763-x
3. Mandalà M, Califano L, Casani AP, et al. Double-Blind Randomized Trial on the Efficacy of the Forced Prolonged Position for Treatment of Lateral Canal Benign Paroxysmal Positional Vertigo. *Laryngoscope.* 2021;131(4):E1296-E1300. doi:10.1002/lary.28981
4. Zuma E Maia F, Ramos BF, Cal R, Brock CM, Mangabeira Albernaz PL, Strupp M. Management of Lateral Semicircular Canal Benign Paroxysmal Positional Vertigo. *Front Neurol.* 2020;11:1040. Published 2020 Sep 15. doi:10.3389/fneur.2020.01040

Summary of Monthly Abstract of the Week Topics From July 2017 to present

Summary of February 2022 Topic: BPPV Updates

The first article investigated the incidence and characteristics of pseudo-spontaneous nystagmus (PSN) in benign paroxysmal positional vertigo involving the lateral canal (LC-BPPV) and evaluated the correlation between PSN and the bow and lean test. In 131 cases of LC-BPPV, the nystagmus in the natural sitting position, supine head roll, Dix-Hallpike, and the bow and lean tests was examined using video-oculography goggles. The subtypes were classified as geotropic or apogeotropic. PSN was found in 25 cases (19.1%) of the LC-BPPV patients with no significant difference in the incidence of PSN in the subtypes. In addition, 21 of the 25 patients exhibited nystagmus consistent with the lean test. In conclusion, PSN can be found in patients with LC-BPPV, irrelevant of subtype; and the pattern of nystagmus was similar to that of the lean test. (1).

The next article was a video case report on canal conversion (canal switching) based on the “vestibular lithiasis” theory. The patient in this article was a 68-year-old female who tested positive for the right posterior canal, right superior canal, and right horizontal canal BPPV within a single session; each successfully treated via the modified Epley, Yacovino, and the Lempert BBQ Roll maneuvers, respectively. Therefore, based on the “vestibular lithiasis” theory, canal switching can be produced by the repositioning maneuvers or by the subsequent control testing maneuvers. (2).

The third article examined the effects of vitamin D and calcium supplementation in preventing BPPV recurrences. A randomized control trial was performed in 1,050 patients with BPPV that were divided into an intervention or observation group. In the intervention group, patients took 400 IU of vitamin D and 500 mg of calcium twice a day for one year when serum vitamin D levels were below the normal range. The observation group had no vitamin D evaluation or supplementation. It was found that the proportion of patients with recurrence of BPPV was lower in the intervention than in the observation group. Therefore, the authors concluded that vitamin D and calcium supplementation may be considered in patients with frequent attacks of BPPV. (3).

The last article evaluated the efficacy of the Li Maneuver as a repositioning maneuver for geotropic horizontal canal benign paroxysmal positional vertigo (HC-BPPV) compared to the Gufoni Maneuver. A retrospective study collected and analyzed 254 cases, of which 87 were treated with Gufoni Maneuver and 167 were treated with Li Maneuver. Statistical analysis showed no significant difference between the two groups. This study concluded that the Li Maneuver is just as effective as the Gufoni Maneuver, but much simpler and faster to perform. (4).

1. Lee, H. J., Ahn, S. K., Yim, C. D., Kim, D. H., & Hur, D. G. (2020). Pseudo-spontaneous nystagmus in lateral semicircular canal benign paroxysmal positional vertigo: Correlation with bow and lean test in a pitch plane. *PLoS One*, 15(11), e0242580.

2. Zanotti, E., & Yacovino, D. A. (2021). Benign Paroxysmal Positional Vertigo: Canal Switching Affecting All Canals During a Single Session. *Otology & Neurotology*, 42(1), e110.

Summary of Monthly Abstract of the Week Topics From July 2017 to present

3. Jeong, S. H., Kim, J. S., Kim, H. J., Choi, J. Y., Koo, J. W., Choi, K. D., & Kim, S. (2020). Prevention of benign paroxysmal positional vertigo with vitamin D supplementation: a randomized trial. *Neurology*, 95(9), e1117-e1125.
4. Zhao, P., Li, J., Ding, Y., Wang, Y., & Zou, S. (2021). Li Maneuver for geotropic horizontal canal benign paroxysmal positional vertigo (HC-BPPV)-A better choice. *American Journal of Otolaryngology*, 103132.

Summary of March 2022 Topic: COVID-19

The March topic for abstract of the month is COVID-19. The first article looked to examine the characteristics of COVID-19 patients especially the developing symptoms of auditory and vestibular difficulties. 300 participants that were diagnosed with COVID-19 participated in the study where common symptoms and complaints were collected during the acute phase and follow up phases. Participants were asked to describe their chief complaints and whether they had specific auditory and vestibular complaints. The results showed that auditory and vestibular symptoms were seen in only a small percentage of patients, but it wasn't their chief complaint, and it was shown that in those who reported the symptoms that the symptoms disappeared in the follow up phases. To conclude, only a small percentage of COVID-19 patients complained of auditory and vestibular symptoms, but it needs to be further researched to determine if the complaints are due to the virus itself, medications, or stress.

(1)

The second article looked to examine the prevalence and characteristics of dizziness and vertigo among patients with COVID-19. Patients went through daily phone calls, nasopharyngeal swab, and were asked about specific symptoms dealing with balance disorders. It was found that symptoms of new-onset dizziness, lightheadedness, disequilibrium, presyncope, and vertigo were seen in the population suggesting that dizziness should be included among the main symptoms of COVID-19. (2)

The third article looked to quantify the amount of balance and proprioception deficit related to post COVID. 66 post COVID-19 patients were analyzed using the robotic device. Hospitalized patients showed the worst performance with dynamic balance activities compared to non-hospitalized patients, but both populations showed bad performance with their eyes closed with dynamic balance activities. Post-COVID patients were significantly more distant from normality with balance activities with their eyes open compared to their eyes closed. This suggests that in post- COVID patients when visual feedback is absent, patients seek more strategies from their proprioception to compensate and improve their performance. This deserves more research for its potential impact on ADLs. (3)

The fourth article was a systematic review examining the literature of audio-vestibular symptoms in COVID-19 patients. From the studies obtained, there were multiple reports of hearing loss, tinnitus, and rotatory vertigo in adults with COVID-19, however there is a low number of high-quality studies comparing the symptoms in COVID-19 compared to controls. (4)

Summary of Monthly Abstract of the Week Topics From July 2017 to present

The fifth article looked at the different experiences of PTs delivering telehealth for vestibular disorders to look at specific barriers, preferences, and concerns. Surveys were given to therapists examining these aspects. Results showed that more than 80% of PTs agreed that telehealth was an effective platform for vestibular PT. When asked whether therapists thought that patients had similar health outcomes with telehealth vs in person care, 68% of therapists agreed. With comfort treating specific diagnoses, more than 50% were comfortable treating posterior or horizontal BPPV compared to more than 75% were comfortable treating peripheral diseases and central disorders. PTs reported several barriers to telehealth from concerns of safety with balance training and challenges with writing a true HEP. To conclude, although there are specific barriers, telehealth for vestibular physical therapy remains to be efficient. (5)

1. Shahidipour Z, Rouhbakhsh N, Tavanai E. Auditory and Vestibular Complaints Among COVID-19 Patients: A Descriptive Survey of 300 Young Patients Over a 3-Month Follow-up Period. *Modern Rehabilitation*. 2022;16(1):69-76. Accessed March 1, 2022.
2. Aldè M, Barozzi S, Di Bernardino F, et al. Prevalence of symptoms in 1512 COVID-19 patients: have dizziness and vertigo been underestimated thus far? [published online ahead of print, 2022 Jan 30]. *Intern Emerg Med*. 2022;1-11. doi:10.1007/s11739-022-02930-0
3. Gervasoni F, LoMauro A, Ricci V, et al. Balance and visual reliance in post-COVID syndrome patients assessed with a robotic system: a multi-sensory integration deficit. *Neurol Sci*. 2022;43(1):85-88. doi:10.1007/s10072-021-05647-8
4. Almufarrij I, Munro KJ. One year on: An updated systematic review of SARS-CoV-2, COVID-19 and audio-vestibular symptoms. *Int J Audiol*. 2021;60(12):935-945. doi:10.1080/14992027.2021.1896793
5. Harrell RG, Schubert MC, Oxborough S, Whitney SL. Vestibular Rehabilitation Telehealth During the SAEA-CoV-2 (COVID-19) Pandemic. *Front Neurol*. 2022;12:781482. Published 2022 Jan 20. doi:10.3389/fneur.2021.781482

Summary of April 2022 Topic: Vestibular Migraine

The first article was a review summarizing the main tests available for diagnosing Vestibular Migraine (VM), including the accuracy of biomarkers for VM diagnosis. All diagnosis records and main VM information published over the past ten years were reviewed. Based on expert opinion, it was concluded that the search for biomarkers will allow a reclassification of VM and improve the precision of a VM diagnosis (1).

The following article assessed kinesiophobia and its association with three clinical factors, fear of falling, dizziness disability, and migraine disability in the migraine population. Eighty patients with migraine were divided into two groups; migraine without kinesiophobia (MoK) and migraine with kinesiophobia (MK), based on the Tampa Scale for Kinesiophobia. In addition, both groups used validated

Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

questionnaires to determine their fear of falling, dizziness disability, and migraine disability. The results showed increased levels of all three clinical factors in the MK group compared to the MoK, concluding that kinesiophobia should be considered in clinical practice when evaluating migraine patients (2).

The third article aimed to compare the relative efficacy of magnesium, metoclopramide, and prochlorperazine for the treatment of migraines in the emergency department (ED). This prospective study analyzed 157 patients who presented to the ED with a diagnosis of migraine from August 2019 to March 2020. These patients were randomized to receive either magnesium sulfate, metoclopramide, or prochlorperazine intravenously. Pain reduction using the Numeric Rating Scale (NRS) after 30 minutes of infusion initiation was the primary outcome. In addition, the ED length of stay, the necessity for rescue analgesia, adverse effects, and the NRS at 60 and 120 minutes were also assessed. Based on the results, there was no significant difference in the change in pain scores at 30 minutes, ED length of stay, rescue analgesia, or adverse effects between all three drugs. The authors concluded that magnesium could be used as an adjunctive agent for the treatment of migraines or as a safe alternative when other agents such as prochlorperazine or metoclopramide are not appropriate (3).

The last article was a review that focused on the effects of various disorders on psychiatric comorbidity and the anxiety associated with vertigo. In addition, it discussed the intimate structural and functional linkage of the vestibular and anxiety systems. Based on several studies reviewed, it was found that there are higher rates of psychiatric comorbidity in patients with vestibular migraines and Meniere's disease and lower rates in patients with vestibular neuritis and bilateral vestibulopathy (BVP). In addition, chronic unilateral and BVP patients reported less anxiety related to vertigo. The authors summarized that the anxiety level ('excess anxiety' or 'less anxiety') of vestibular dysfunctions may be based on whether there is excitation or loss of vestibular function (4).

1. Perez-Carpena P, Lopez-Escamez JA. Do we need to reconsider the classification of vestibular migraine?. *Expert Rev Neurother*. 2021;21(5):503-516. doi:10.1080/14737175.2021.1908129
2. Pinheiro CF, Bevilaqua-Grossi D, Florencio LL, et al. Is kinesiophobia related to fear of falling, dizziness disability, and migraine disability in patients with migraine? [published online ahead of print, 2021 Oct 27]. *Physiother Theory Pract*. 2021;1-9. doi:10.1080/09593985.2021.1996496 PMID: 34704520 DOI: 10.1080/09593985.2021.1996496
3. Kandil M, Jaber S, Desai D, et al. MAGraïne: Magnesium compared to conventional therapy for treatment of migraines. *Am J Emerg Med*. 2021;39:28-33. doi:10.1016/j.ajem.2020.09.033
4. Brandt T, Dieterich M. 'Excess anxiety' and 'less anxiety': both depend on vestibular function. *Curr Opin Neurol*. 2020;33(1):136-141. doi:10.1097/WCO.0000000000000771

Summary of May 2022 Topic: BPPV Diagnosis and Treatment Updates

The May topic for abstract of the month is BPPV. The first article looked to examine the effect of ocular fixation on positional nystagmus in BPPV patients. Positional maneuvers were performed on 58 subjects

Summary of Monthly Abstract of the Week Topics From July 2017 to present

who had BPPV where ocular fixation was then initiated, and short phase velocity was recorded. It was found that ocular fixation significantly suppressed positional nystagmus in BPPV patients during positional maneuvers. (1)

The second article looked to reveal the effectiveness of the newly Gans repositioning maneuver for treatment of posterior canal BPPV patients. This article conducted a systematic review of RCTs where the Gans maneuver was identified and showed the proportion of patients who improved because of the maneuver. It was shown that improvement was seen in almost all patients with the Gans maneuver in three trials with no significant side effects from the treatment showing that the Gans maneuver is a safe and effective treatment for patients with posterior canal BPPV. (2)

The third article looked to determine whether a single session treatment of unilateral posterior canal BPPV is more effective as a treatment recommendation compared to conventional treatment. RCT was completed consisting of 53 patients diagnosed with unilateral posterior canal BPPV where 26 of them were assigned to the treatment session, while the other 27 were assigned to weekly treatment. It was found that for those in the single session group; all outcome measures were significantly lower including average and total cost of care and loss of productivity for the company due to patients' medical visits. To conclude, the single session was proven to be faster, more efficient, and able to reduce direct and indirect cost of care related to unilateral posterior canal BPPV. (3)

The last article looked to explain BPPV with its different types and treatment. BPPV is characterized by positional vertigo where there are two types which are canalolithiasis and cupulolithiasis. In the diagnosis of BPPV, the characteristics of vertigo, triggered movements of vertigo, and duration of vertigo are all important to diagnose. Dix-Hallpike test is a positional nystagmus test used to diagnose lateral canal BPPV. Canalith repositioning procedures are done to treat BPPV with Epley maneuver being done for posterior canal BPPV, while the Gufoni maneuver is done for lateral canal BPPV. (4)

1. Özel HE, Karakuzu AT, Temir H, Alpay M, Özdoğan F, Genç S. Effect of ocular fixation on positional nystagmus in BPPV patients [published online ahead of print, 2022 Apr

19]. *Int J Audiol.* 2022;1-6. doi:10.1080/14992027.2022.2062579

2. Joshi D, Gyanpuri V, Pathak A, et al. Gans repositioning maneuver for the posterior canal BPPV patients: systematic review and meta-analysis [published online ahead of print, 2022 Apr 23]. *Eur Arch Otorhinolaryngol.* 2022;10.1007/s00405-022-07396-6.

doi:10.1007/s00405-022-07396-6

3. Franco-Gutiérrez V, Pérez-Guillén V, Gil-Aguilar MT, et al. Comparative analysis of the efficiency of two treatment protocols for posterior canal benign paroxysmal positional vertigo [published online ahead of print, 2021 Jan 19]. Análisis comparativo de la eficiencia de dos protocolos de tratamiento del vértigo posicional paroxístico benigno del conducto semicircular posterior [published online ahead of print, 2021 Jan 19]. *Acta Otorrinolaringol Esp (Engl Ed).* 2021;S0001-6519(20)30201-6.

doi:10.1016/j.otorri.2020.11.006

Summary of Monthly Abstract of the Week Topics From July 2017 to present

4. Auris Nasus Larynx. 2022 Apr 3:S0385-8146(22)00066-9. doi: 10.1016/j.anl.2022.03.012.

Summary of June 2022 Topic: Meniere's Disease

The first article presented was a Clinical Practice Guideline (CPG) on Meniere's Disease (MD). The purpose of the CPG was to use the best available published evidence to evaluate the treatment approaches for MD and report on their efficacy in controlling MD symptoms. The CPG consisted of 16 evidence-based action statements, including two statements that defined the role of vestibular rehabilitation (VR). It concluded that VR should be offered to patients with MD, who experienced chronic imbalance, and should not be recommended in the acute management of MD related vertigo attacks. (1)

The second article was a retrospective analysis that aimed to investigate the relationship (concordance or discordance) of caloric testing and video head impulse test (vHIT) results in patients with Meniere's Disease (MD). Patients were placed in three groups after undergoing both caloric testing and vHIT: normal, concordant, and discordant. A total of 644 patients were examined, but only 73 were diagnosed with definite MD. The results showed the patient's with MD had a greater discordance of results with the caloric testing being asymmetrical and vHIT normal. Therefore, the authors concluded that findings of asymmetric caloric function with a normal vHIT may function as a diagnostic marker for MD. (2)

The third article was a systematic review and meta-analysis that aimed to examine the frequency of BPPV in patients with Meniere's Disease (MD) and its clinical features. A total of 20 studies were eligible and included in the systemic review. Results showed an average frequency of 14% of BPPV in MD (affecting females more than males); 2.5 times greater frequency of BPPV when patients were examined more than once; and the horizontal semicircular canal was more commonly affected. In conclusion, BPPV in patients with MD have a high frequency and recurrence rate that may require more BPPV treatments. (3)

The last article was a retrospective study assessing how useful Skull Vibration-Induced Nystagmus Test (SVINT) can be to differentiate between Vestibular Migraines (VM) and Meniere's Disease (MD). This study compared 605 patients diagnosed with MD with 200 patients diagnosed with VM. While the patient was seated, the SVINT was performed at a frequency of 100Hz perpendicular to the skin over the mastoid process with a force around 1kg for a total of three trials on each patient's mastoid. If horizontal nystagmus towards the unaffected side was elicited in all six trials, the test was considered positive. The SVINT was positive in 59.2% of patients with MD and only 6% in patients with VM. The authors concluded that the results support previous findings of the SVINT being positive in peripheral vestibular disorders and the different pathophysiological mechanisms of vertigo in VM likely reflects the low rate of positivity with SVINT. (4)

1. Basura, G. J., Adams, M. E., Monfared, A., Schwartz, S. R., Antonelli, P. J., Burkard, R., & Buchanan, E.M. (2020). Clinical practice guideline: Ménière's disease. *Otolaryngology–Head and Neck Surgery*, 162(2_suppl), S1-S55.

Summary of Monthly Abstract of the Week Topics From July 2017 to present

2. Hannigan, I. P., Welgampola, M. S., & Watson, S. R. (2021). Dissociation of caloric and head impulse tests: a marker of Meniere's disease. *Journal of Neurology*, 268(2), 431-439.
3. Kutlubaev, M. A., Xu, Y., & Hornibrook, J. (2021). Benign paroxysmal positional vertigo in Meniere's disease: systematic review and meta-analysis of frequency and clinical characteristics. *Journal of Neurology*, 268(5), 1608-1614.
4. Teggi, R., Gatti, O., Familiari, M., Cangiano, I., & Bussi, M. (2021). Skull Vibration-Induced Nystagmus Test (SVINT) in Vestibular Migraine and Menière's Disease. *Audiology Research*, 11(4), 603-608.

Summary of July 2022 Topic: Pediatrics and Vestibular Disorders

The first article expresses how arduous vertigo can be to evaluate in young children. Despite vertigo being a very common pediatric emergency that isn't dangerous in nature, it can disguise other serious illnesses which can be dangerous. A study was conducted over a ten-year period comprised of 757 children in the pediatric emergency department for vertigo. It was concluded that children younger than six years old have a higher possibility of a CNS disease cause by vertigo than older groups of children. (1)

The second article explains that according to the ICVD and the Migraine Classification Subgroup of the International Headache Society. Recurrent vertigo has the potential to lead to childhood migraines and there are three types of disorders. These three types are vestibular migraine of childhood, probable vestibular migraine of childhood and recurrent vertigo of childhood, which have to occur under the age of 18 years old and satisfy the diagnostic criteria. In order to expand research on childhood migraines and its three disorders it is recommended that recurrent vertigo of childhood be studied further so that subtypes of the three disorders can be explored and analyzed. (2)

The third article covers a study done over an eight-month period that seeks to discover audio-vestibular complaints within COVID-19 patients and determine if gender and age have an affect on their symptoms. The study was comprised of 1,437 patients and 524 of them complained of more frequent audio vestibular symptoms. 52 patients experienced the more common COVID-19 symptoms such as loss of taste/smell and intermittent vertigo. In conclusion, audio-vestibular symptoms are more commonly experienced at the later part of the 14-day sickness period. (3)

References

1. Grasso, A., Poropat, F., Kamagni Vodié, T., Ghirardo, S., & Barbi, E. (2020). How age matters in the assessment of vertigo in the pediatric emergency department. *Pediatric Emergency Care*, 38(1), e43-e46. <https://doi.org/10.1097/pec.0000000000002242>
2. Van de Berg, R., Widdershoven, J., Bisdorff, A., Evers, S., Wiener-Vacher, S., Cushing, S. L., Mack, K. J., Kim, J. S., Jahn, K., Strupp, M., & Lempert, T. (2021). Vestibular migraine and recurrent vertigo of childhood: Diagnostic criteria consensus document of the classification committee of vestibular

Summary of Monthly Abstract of the Week Topics From July 2017 to present

disorders of the Bárány society and the International Headache Society. *Journal of Vestibular Research*, 31(1), 1-9. <https://doi.org/10.3233/ves-200003>

3. ÇILDIR, B. (2021). An age-related investigation of audio-vestibular symptoms in individuals with COVID-19 positive. *Cukurova Medical Journal*, 46(4), 1537-1547. <https://doi.org/10.17826/cumj.987374>

Summary of August 2022 Topic: Vestibular Paroxysmia

The first article aimed to review the existing literature on Vestibular Paroxysmia (a neurovascular compression syndrome) and to determine the most appropriate imaging technique for this syndrome. Studies were found via a PubMed database search with findings that 3D Constructive Interference in Steady-State (CISS) MRI sequence was the most reliable to image the neurovascular contact between the eighth cranial nerve and a vessel (1).

The purpose of the second article was to describe the clinical symptoms and laboratory vestibular findings in patients with Vestibular Paroxysmia (VP) and to evaluate the long-term clinical course over the years. Seventy-three patients were identified as having probable VP (pVP) and 73 as definite VP (dVP) from a tertiary dizziness center registry. Clinical symptoms included the frequency of attacks from 5 to 30 per day, with two-thirds of those attacks occurring spontaneously and one-quarter triggered by head movements. In addition, the most frequent type of vertigo was spinning in dVP, and 70% of patients reported no accompanying symptoms. Laboratory vestibular findings included evidence of hyperventilation-induced nystagmus, a three and six-degree deviation in Subjective Visual Vertical, and no evidence of complete loss of peripheral vestibular function. The long-term prognosis of VP appears favorable for three-quarters of patients remaining attack-free in dVP and half without needing medication; the frequency was significantly reduced for those with ongoing attacks (2).

The third article was a cross-sectional observational study that explored the long-term course of outcomes in Vestibular Paroxysmia (VP). Seventy-three patients diagnosed with definite or probable VP were included in the study; however, only 44 were part of the follow-up. Patients were contacted by telephone to complete a study-specific questionnaire; their baseline data was retrospectively collected from electronic medical records. At follow-up (mean follow-up time was 3.4 years), 72% of patients still experienced vertigo attacks with an attack frequency of 153 in the past six months, 19.7% reported using or have used carbamazepine, and 71% reported having limitations in one or more health-related quality of life (HRQoL). The authors concluded an unfavorable prognosis in patients with VP per vertigo attacks and HRQoL limitations (3).

The final article compared the clinical, electrophysiological, radiological, and therapeutic outcomes of pediatric patients diagnosed with probable Vestibular Paroxysmia. A total of 16 pediatric patients were included in the retrospective descriptive comparative study and divided into those with narrowed internal auditory canal (IAC) and those with neurovascular cross-compression syndrome (NVCC). Findings showed similar auditory complaints, hearing, vestibular, and electrophysiological status in both groups, with a narrowed IAC more commonly present in adolescence. The study suggested carefully

Summary of Monthly Abstract of the Week Topics From July 2017 to present

assessing the IAC diameter in the coronal plane and the axial plane. Lastly, it was found that treatment with sodium-channel blockers drugs can help relieve vertigo and normalize electrophysiological findings. The authors concluded that VP could originate from a narrowed IAC (4).

1. Karamitros A, Kalamatianos T, Stranjalis G, Anagnostou E. Vestibular paroxysmia: Clinical features and imaging findings; a literature review. *J Neuroradiol.* 2022;49(2):225-233. doi:10.1016/j.neurad.2021.07.007
2. Steinmetz K, Becker-Bense S, Strobl R, Grill E, Seelos K, Huppert D. Vestibular paroxysmia: clinical characteristics and long-term course [published online ahead of print, 2022 May 20]. *J Neurol.* 2022;10.1007/s00415-022-11151-6.
3. Hanskamp LAJ, Schermer TR, van Leeuwen RB. Long-term Prognosis of Vertigo Attacks and Health-related Quality of Life Limitations in Patients With Vestibular Paroxysmia. *Otol Neurotol.* 2022;43(4):e475-e481. doi:10.1097/MAO.0000000000003465
4. Idriss SA, Thai-Van H, Altaisan R, Ltaief-Boudrigua A, Reynard P, Ionescu EC. The Narrowed Internal Auditory Canal: A Distinct Etiology of Pediatric Vestibular Paroxysmia. *J Clin Med.* 2022;11(15):4300. Published 2022 Jul 25. doi:10.3390/jcm11154300

Summary of September 2022 Topic: Acute Dizziness in the ED

The September topic for the abstract of the month is Acute Dizziness in the ED. The first article was a systematic review and meta-analysis examining whether ED physicians could perform the HINTS examination or STANDING algorithm to differentiate between central vs non- central vertigo and compare it to the findings of more specialized physicians. Previously, research has shown that ED physicians are unable to use the HINTS examination without having the appropriate training. From the analysis, there were 2 studies of 450 patients examined by 11 emergency physicians. The analysis showed that ED physicians who had received prior training were able to utilize the STANDING algorithm with a sensitivity of .96 and a specificity of .88, but there was no data collected for the HINTS examination. To conclude, it was found that when ED physicians are trained that they can use the STANDING algorithm with confidence. (1)

The second article compared characteristics of acute nystagmus in different diagnoses such as posterior circulation stroke and acute vestibular neuritis in the ER within 24 hours of presentation. The study utilized ER-based video nystagmography to record nystagmus in 101 patients with posterior circulation stroke and 104 patients with acute vestibular neuritis. For patients with posterior circulation stroke, results showed that 44 out of 101 patients had no spontaneous nystagmus, while the rest had primary position horizontal, vertical, and torsional nystagmus. It was found that the patients with horizontal nystagmus had unidirectional “peripheral appearing” nystagmus, while others had gaze-evoked nystagmus. In comparison, 102 out of the 104 patients with acute vestibular neuritis had primary position horizontal nystagmus, while none had gaze-evoked nystagmus. From this, it was concluded that

Summary of Monthly Abstract of the Week Topics From July 2017 to present

nystagmus is always present in acute vestibular neuritis, while unidirectional peripheral appearing horizontal nystagmus is seen in posterior circulation stroke. (2)

The third article was a systematic review and meta-analysis examining the diagnostic test accuracy of neuroimaging for patients with acute vertigo or dizziness. It was found that non-contrast CT had very low sensitivity, while MRI will miss 1 in 5 patients with stroke if imaging is obtained early after symptom onset. To conclude, the evidence does not support neuroimaging as the only tool for ruling out stroke and other causes in patients with dizziness and vertigo presenting to the ED. (3)

The last article was a systematic literature review looking to examine the pathologies of neurological vertigo in childhood and adulthood to help clinicians to a proper diagnosis. From this, the review aimed to develop an algorithm to help ER physicians approach patients with isolated vertigo. (4)

1. Nakatsuka M, Molloy EE. The HINTS examination and STANDING algorithm in acute vestibular syndrome: A systematic review and meta-analysis involving frontline point-of-care emergency physicians. PLoS One. 2022;17(5):e0266252. Published 2022 May 5. doi:10.1371/journal.pone.0266252
2. Nham B, Akdal G, Young AS, et al. Capturing nystagmus in the emergency room: posterior circulation stroke versus acute vestibular neuritis [published online ahead of print, 2022 Jul 18]. J Neurol. 2022;10.1007/s00415-022-11202-y. doi:10.1007/s00415-022-11202-y
3. Shah VP, Oliveira J E Silva L, Farah W, et al. Diagnostic accuracy of neuroimaging in emergency department patients with acute vertigo or dizziness: A systematic review and meta-analysis for the guidelines for reasonable and appropriate care in the emergency department [published online ahead of print, 2022 Jul 25]. Acad Emerg Med.2022;10.1111/acem.14561. doi:10.1111/acem.14561
4. Pellegrino N, Di Stefano V, Rotondo E, et al. Neurological vertigo in the emergency room in pediatric and adult age: systematic literature review and proposal for a diagnostic algorithm. Ital J Pediatr. 2022;48(1):125. Published 2022 Jul 27. doi:10.1186/s13052-022-01313-7

Summary of October 2022 Topic: Rehabilitation Implications of Vestibular Surgical Interventions

The month of October focused on surgical interventions and their implications on a vestibular patient's postural control, gait, quality of life and hearing. Highlighted interventions include the vestibular and cochlear implants, gamma knife and neurectomy.

The first article measured the clinical effects in patients with bilateral vestibular hypofunction after a unilateral vestibular implant. The clinical outcomes of eight patients were analyzed via the Bruininks-Oseretsky Test, modified Romberg, Dynamic Gait Index, gait speed, Timed Up and Go, pure-tone auditory detection thresholds, speech discrimination scores, and quality of life. The patients' results from pre-implantation were compared with the results at six months and one year, with the device set

Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

in treatment and placebo mode. All clinical outcomes of posture, gait, and quality of life showed improvement at six months and the one year mark, however, ipsilateral hearing reduction occurred in seven patients. (1)

The second article aimed to analyze whether there was a change of vestibular function in patients following cochlear implant surgery. This systematic review with meta-analysis included databases between 1980 and 2021 and found 38 articles that met their inclusion criteria. Results indicated a statistically significant difference between vestibular-evoked myogenic potential (VEMP) and caloric tests after cochlear implant surgery; however, VEMP was the only test that evaluated the vestibular apparatus as having abnormal results. (2)

The final article investigated the changes in peripheral vestibular function and central compensation by measuring head motion kinematics in patients with vestibular neurectomies due to vestibular schwannoma. A small head-mounted sensor quantified data during the Functional Gait Assessment (FGA). Measurements were made before surgery followed by six weeks and six months post surgery. Results showed a significant difference in kinematics between normal subjects and those with vestibular schwannoma (both pre-surgery and post-surgery). Kinematics at six weeks post-operative were significantly altered when compared to pre-operative. There was minimal change between six weeks and six months post-operative. The authors concluded that head movement kinematics provide valuable information about vestibular damage and compensation, and speculate changes in head motion strategy prior to surgery may affect long-term vestibular compensation. (3)

1. Chow, M. R., Ayiotis, A. I., Schoo, D. P., Gimmon, Y., Lane, K. E., Morris, B. J., & Della Santina, C. C. (2021). Posture, gait, quality of life, and hearing with a vestibular implant. *New England Journal of Medicine*, 384(6), 521-532.
2. Vaz, F., Petrus, L., Martins, W., Silva, I., Lima, J., Santos, N., & Bahmad, F. (2022). The effect of cochlear implant surgery on vestibular function in adults: A meta-analysis study. *Frontiers in neurology*, 13.
3. Zobeiri, O. A., Mischler, G. M., King, S. A., Lewis, R. F., & Cullen, K. E. (2021). Effects of vestibular neurectomy and neural compensation on head movements in patients undergoing vestibular schwannoma resection. *Scientific reports*, 11(1), 1-14.

Summary of November 2022 Topic: Vestibular Disorders

The November topic for abstract of the month is Vestibular Disorders. The first article looked to discuss the association between vestibular neuritis and COVID-19 vaccination. COVID-19 is a severe respiratory syndrome and presents with pneumonia as the most serious manifestation. As a result, COVID-19 vaccination has been an important interest globally due to the rapid expansion of COVID-19 with its high rates of mortality and morbidity. On the other hand, vestibular neuritis is an acute vestibular syndrome that causes spontaneous vertigo due to unilateral vestibular difficulties leading to nausea and unsteadiness that can last for weeks. From this study, there was one individual who was discovered to

Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

develop vestibular neuritis within 3 days after COVID-19 vaccination, but there have been no further reports of the diagnosis of vestibular neuritis after COVID-19 vaccination, and the association is unclear, but clinicians should be aware that vestibular neuritis may occur as an adverse event of COVID-19 vaccination. (1)

The second article looked to describe and discuss the topic of vestibular asthenopia. Vestibular asthenopia is a sensory discomfort that consist of multiple subjective symptoms specifically visual that has originated from a transient vestibular incident. It can happen because of a sudden global central disorder following a chain of events in response to an initial vestibular disease. It results in inadequate processing and difficulty integrating afferent visual and vestibular input leading to ocular fatigue, pain with eye movement, and sensitivity to retinal slip. (2)

The third article looked to discuss and explore the topic of vestibular migraines and their recent research developments. The review discusses how patients diagnosed with vestibular migraines experience persistent vestibular symptoms, even in the absence of acute attacks, which in the long run significantly impact overall quality of life. The review further discusses how current and previous studies support the benefit of migraine treatments to be effective for vestibular migraines, including neuromodulation and calcitonin gene-related peptide monoclonal antibodies. Vestibular migraines affect almost 3% of the population and remains under diagnosed. With new and recent research, diagnostic criteria can help clinicians diagnose vestibular migraines more accurately in adults and children. (3)

The last article was a narrative review looking to discuss the current diagnosis and treatment of vestibular neuritis. For the diagnosis of vestibular neuritis, the key features include “abrupt true-whirling vertigo lasting for more than 24 hours and no presence of cochlear symptoms or other neurological symptoms”. There are various diagnostic tests such as the head impulse test, bithermal caloric test, and vestibular evoked myogenic potential test are all options utilized to diagnose. There are a variety of treatments for vestibular neuritis such as symptomatic therapy, drug therapy, and vestibular rehabilitation therapy. (4)

1. Jeong J. Vestibular neuritis after COVID-19 vaccination. *Hum Vaccin Immunother.* 2021;17(12):5126-5128. doi:10.1080/21645515.2021.2013085

2. Bauwens A, Larock F. Vestibular asthenopia. *J Fr Ophtalmol.* 2021;44(10):1560-1565. doi:10.1016/j.jfo.2021.05.008

3. Beh SC. Vestibular Migraine. *Curr Neurol Neurosci Rep.* 2022;22(10):601-609. doi:10.1007/s11910-022-01222-6

4. Bae CH, Na HG, Choi YS. Current diagnosis and treatment of vestibular neuritis: a narrative review. *J Yeungnam Med Sci.* 2022;39(2):81-88. doi:10.12701/yujm.2021.01228

Summary of Monthly Abstract of the Week Topics From July 2017 to present

Summary of December 2022 Topic: Red Flags

The purpose of the first article was to determine if the presence of vertigo was a red flag for brain metastasis (BM) in patients with cancer. The study reviewed the neuro-oncologic consultation of patients with confirmed systemic cancer. Of 3220 patients, 22.5% who experienced vertigo were diagnosed with BM; vertigo was the initial symptom in 6% of patients with BM. In addition, there was a higher risk of having BM in patients presenting with vertigo and accompanied headache, ataxia, seizures, visual symptoms, speech impairments, altered mental status, focal weakness, or focal sensitive complaints. Therefore, the study concluded that vertigo and these accompanying symptoms should be considered red flags in patients with systemic cancer. (1)

The second article provided an overview of rare vestibular disorders to help determine efficient diagnosis and treatment. The vestibular syndromes were classified as acute, episodic, and chronic. Examples of disorders include Tumarkin attacks, third window syndromes, endolymphatic sac tumors, rare BPPV variants, and autoimmune inner ear diseases. The article discusses the pathophysiology, possible causes, clinical presentation, diagnostic approach, and available therapies/treatments. In addition, it provides red flags for potentially dangerous conditions (e.g., labyrinthine infarction/hemorrhage). (2)

The next article aimed to identify the characteristics and red flags present with acute nystagmus (AN) in pediatric patients. The authors retrospectively reviewed the clinical records of 206 patients between 0 to 18 years old who presented with AN in the emergency department. The study found that migraine and vestibular disorders were the most common causes of AN. In addition, brain tumors were a common underlying urgent condition (UC) in 8.3% of all cases. Red flags for underlying UC included cranial nerve palsy, ataxia, and strabismus. Therefore, the authors concluded that AN should be considered an alarming finding in the pediatric population. (3)

The final article discusses clinical features and red flags that should be considered to help distinguish central lesions from benign paroxysmal positional vertigo (BPPV). Seven cases were presented that had an initial diagnosis of BPPV but then found to have been from a central nature. The case series suggests that isolated downbeat nystagmus and apogeotropic horizontal nystagmus (on supine roll test) should be considered additional red flags. (4)

1. Cacho-Díaz, B., Salmerón-Moreno, K., Mendoza-Olivas, L. G., Reynoso-Noverón, N., & Gómez-Amador, J. L. (2019). Vertigo in patients with cancer: Red flag symptoms. *Journal of clinical neuroscience : official journal of the Neurosurgical Society of Australasia*, 69, 175–178. <https://doi.org/10.1016/j.jocn.2019.07.071>

2. Długaiczek J. (2021). Rare Disorders of the Vestibular Labyrinth: of Zebras, Chameleons and Wolves in Sheep's Clothing. *Seltene Erkrankungen des vestibulären Labyrinths: von Zebras, Chamäleons und Wölfen im Schafspelz*. *Laryngo- rhino- otologie*, 100(S 01), S1–S40. <https://doi.org/10.1055/a-1349-7475>

Summary of Monthly Abstract of the Week Topics From July 2017 to present

3. Garone, G., Suppiej, A., Vanacore, N., La Penna, F., Parisi, P., Calistri, L., Palmieri, A., Verrotti, A., Poletto, E., Rossetti, A., Cordelli, D. M., Velardita, M., d'Alonzo, R., De Liso, P., Gioè, D., Marin, M., Zagaroli, L., Grosso, S., Bonfatti, R., Mencaroni, E., ... Raucci, U. (2020). Characteristics of Acute Nystagmus in the Pediatric Emergency Department. *PPediatrics*, 146(2), e20200484.
<https://doi.org/10.1542/peds.2020-0484>
4. Joshi, P., Mossman, S., Luis, L., & Luxon, L. M. (2020). Central mimics of benign paroxysmal positional vertigo: an illustrative case series. *Neurological sciences : official journal of the Italian Neurological Society and of the Italian Society of Clinical Neurophysiology*, 41(2), 263–269.
<https://doi.org/10.1007/s10072-019-04101-0>

Summary of January 2023 Topic: BPPV

The January topic for abstract of the month was the topic of BPPV. The first article looked to observe the horizontal, vertical, and torsional characteristics of nystagmus in patients with posterior semicircular canal BPPV and look at its correlation with Ewald's. There were 84 patients enrolled with BPPV where the latency, duration, direction, and slow-phase velocity induced by the Dix Hallpike test in the head-hanging and sitting positions were recorded. It was found that the fast phase of the vertical components and torsional components of left and right ear were upward, clockwise, and counterclockwise, and the horizontal components were mainly contralateral. When the patients were returned to the seated position, the fast phase of the vertical and torsional components of nystagmus was reversed. Only 54 patients had horizontal components of nystagmus, and 32 of them remained in the same direction. It was concluded that from the three components of nystagmus induced by the Dix Hallpike test that the vertical component was the strongest, and the horizontal component was the weakest. (1)

The second article looked to investigate the occurrence of canal switch- benign paroxysmal positional vertigo (CS-BPPV), the characteristics, and underlying mechanisms. Data was assessed from 1,099 patients with single-canal BPPV where patients were divided into CS-BPPV and non-CS BPPV where the baseline characteristics and nystagmus characteristics were compared between the two groups. It was found that of these patients that 51/1,099 developed CS-BPPV. There were no statistically significant differences between the two groups. The intensity of nystagmus and the incidence of direction-reversing nystagmus was significantly greater in the CS-BPPV group compared to the non-CS-BPPV. To conclude, CS in BPPV is uncommon, and it was found that patients with anterior canal BPPV are more likely to develop CS. The occurrence of CS-BPPV may be related to the anatomical structure of the semicircular canals showing that when the canals contain large accumulations of otolith particles that CS may be more common. (2)

The third article looked to investigate the function of otoliths in patients with BPPV before, after, and 1 month after repositioning and look at the possible compensation mechanisms. There were 36 patients with confirmed BPPV and 36 healthy patients where the virtual reality static subjective visual vertical and subjective visual horizontal were measured for the healthy group at inclusion, and for the BPPV group, measured before, after, and 1 month after repositioning. It was found that before repositioning

Summary of Monthly Abstract of the Week Topics From July 2017 to present

that the otolithic organ function of BPPV patients was impaired with no difference between the healthy and affected ear, while after repositioning, there was a transient recovery of otolithic dysfunction followed by a decline to similar levels to before repositioning. (3)

The last article looked to compare the efficacy of a home-based repositioning procedure compared to the standard Epley's maneuver in treating patients with posterior canal BPPV. Patients were randomized into two groups where one group received the standard treatment and the other received the home based. The vertigo scale, duration of nystagmus during Dix Hallpike, and frequency of vertigo were documented on visits and compared between both groups. It was found that there was no significant difference in the baseline characteristics in both groups; both groups had significant improvement of symptoms at the end of the study. To conclude, the home-based repositioning is an effective procedure for the treatment of patients with posterior canal BPPV. (4)

1. Liu, Y., Zhang, X., Deng, Q., Liu, Q., Wen, C., Wang, W., & Chen, T. (2022). The 3D characteristics of nystagmus in posterior semicircular canal benign paroxysmal positional vertigo. *Frontiers in neuroscience*, 16, 988733. <https://doi.org/10.3389/fnins.2022.988733>
2. Wu, Y., Song, N., Ling, X., Li, X., Feng, Y., Xing, Y., Gu, P., & Yang, X. (2022). Canal switch in benign paroxysmal positional vertigo: Clinical characteristics and possible mechanisms. *Frontiers in neurology*, 13, 1049828. <https://doi.org/10.3389/fneur.2022.1049828>
3. Zhao, C., Yang, Q., & Song, J. (2022). Dynamic changes of otolith organ function before and after repositioning in patients with benign paroxysmal positional vertigo detected by virtual reality auxiliary technology: A cohort study. *Frontiers in neurology*, 13, 1007992. <https://doi.org/10.3389/fneur.2022.1007992>
4. Ranju, R. L., Lepcha, A., Mammen, M. D., Vasanthan, L. T., Augustine, A. M., & Philip, A. (2022). An Effective Home-Based Particle Repositioning Procedure for Posterior Canal Benign Paroxysmal Positional Vertigo (BPPV). *Indian journal of otolaryngology and head and neck surgery : official publication of the Association of Otolaryngologists of India*, 74(4), 516–523. <https://doi.org/10.1007/s12070-021-03021-w>

Summary of February 2023 Topic: Coexistence of Vestibular Deficits and Psychiatric Disorders

For the month of February, we take an updated look at the coexistence of vestibular deficits and psychiatric disorders.

The first article examined observed postural sway and perceived instability in patients with Persistent Postural Perceptual Dizziness (PPPD). It included 19 patients with PPPD, 10 healthy controls, and 10 with bilateral vestibulopathy (BV). Observed sway was measured on a force plate while the participants completed quiet standing on a firm surface with eyes closed; perceived instability was verbally rated using a 0-10 ranked scale; and quantified perceived instability (reproduced sway) was measured on the force plate with eyes open. Results showed greater perceived instability and reproduced sway in PPPD



Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

patients compared to observed sway; in patients with BV, the observed sway was proportionate to reproduced sway and perceived instability. Therefore, the authors concluded that future biomarkers of PPPD may need to rely less on observed postural sway and rather include measures of perceptual impairment (1).

The second article was a case report of a 40-year-old woman diagnosed with Persistent Postural-Perceptual Dizziness (PPPD) and post-stroke depression (PSD) after an insular stroke. A few days after her stroke, the patient developed chronic dizziness (exacerbated by self-motion, visual motion, and precise visual focus) and depressive symptoms. Her dizziness persisted for another six months despite a normal neurological and vestibular examination. Due to the patient's development of PPPD and PSD occurring at the same time, the authors propose the mechanism of these symptoms was likely due to the function of the insula in spatial orientation and mood regulation and its multiple networks in the human brain (2).

1. Murillo, E. S. P., Bancroft, M. J., Koochi, N., Castro, P., & Kaski, D. (2023). Postural misperception: a biomarker for persistent postural perceptual dizziness. *Journal of Neurology, Neurosurgery & Psychiatry*, 94(2), 165-166. PMID: 35995549 DOI: 10.1136/jnnp-2022-329321
2. Fipps, D. C., Staab, J. P., & Allen, N. D. (2022). Case Report: Right Insular Stroke Causing Simultaneous Onset of a Functional Vestibular Disorder and Psychiatric Disorder—Persistent Postural-Perceptual Dizziness and Post-stroke Depression. *Frontiers in Psychiatry*, 13. PMID: 35530020 DOI: 10.3389/fpsyt.2022.841072

Summary of March 2023 Topic: Vestibular Migraine

The March topic for abstract of the month was the topic of Vestibular Migraine. The first article was a review that looked to explore the current diagnostic criteria for vestibular migraine in the adult and pediatric populations. Studies have revealed new insights, including the wide range of vestibular symptoms, symptoms in the attack-free period, and triggers. There are a plethora of randomized controlled trials finding neuromodulation and calcitonin gene-related peptide monoclonal antibodies as effective treatments for vestibular migraines. With new diagnostic criteria, it can help clinicians diagnose vestibular migraines in adults and children. (1)

The second article looked to discuss the topic of vestibular migraine as a whole. It is a neurological condition that causes episodic vertigo and is thought to be the most common cause of spontaneous episodic vertigo. There has been continued discussion of preventative treatments, but studies have not been able to establish a right treatment due to low quality of evidence. (2)

The third article looked at the impact of vestibular rehabilitation therapy on the treatment of vestibular migraine and looked at the relationship between TBI and vestibular migraine. It was found that vestibular rehabilitation therapy significantly improved self-reported dizziness on the Dizziness Handicap

Summary of Monthly Abstract of the Week Topics From July 2017 to present

Inventory and GAIT assessment in individuals with vestibular migraine. It was also found that a history of TBI impacted outcome on the Dizziness Handicap Inventory. To conclude, vestibular rehabilitation therapy can be seen as an effective treatment option to reduce dizziness and the risk of falls in individuals with vestibular migraine. (3)

The fourth article looked to differentiate vestibular migraine from Meniere's disease. Meniere's disease is classified as a triad of symptoms: recurrent vertigo, fluctuating hearing loss, and tinnitus. It is critical to be able to differentiate the two due to the difference of treatment. Symptoms of vestibular migraine consist of headache, vestibular symptoms, and cochlear symptoms. There are new advances in examination technologies to help differentiate the two. (4)

The last article looked to understand the cognitive dysfunction associated with vestibular migraine. It was found that among patients with vestibular migraine that 40% reported "some" and 12% reported "a lot" of difficulty thinking compared to lower percentages for patients without vestibular migraines. Individuals with vestibular migraines also had increased odds of falls and mobility issues compared to those without. To conclude, it was found that vestibular migraine is associated with cognitive dysfunction, mobility issues, and fall risk. (5)

1. Beh SC. Vestibular Migraine. *Curr Neurol Neurosci Rep.* 2022 Oct;22(10):601-609. doi:10.1007/s11910-022-01222-6. Epub 2022 Aug 31. PMID: 36044103.
2. Smyth D, Britton Z, Murdin L, Arshad Q, Kaski D. Vestibular migraine treatment: a comprehensive practical review. *Brain.* 2022;145(11):3741-3754. doi:10.1093/brain/awac264
3. Stancel-Lewis J, Lau JW, Male A, et al. Vestibular Rehabilitation Therapy for the Treatment of Vestibular Migraine, and the Impact of Traumatic Brain Injury on Outcome: A Retrospective Study. *Otol Neurotol.* 2022;43(3):359-367. doi:10.1097/MAO.0000000000003452
4. Chen JY, Guo ZQ, Wang J, et al. Vestibular migraine or Meniere's disease: a diagnostic dilemma [published online ahead of print, 2022 Dec 23]. *J Neurol.* 2022;10.1007/s00415-022-11532-x. doi:10.1007/s00415-022-11532-x
5. Preysner TA, Gardi AZ, Ahmad S, Sharon JD. Vestibular Migraine: Cognitive Dysfunction, Mobility, Falls. *Otol Neurotol.* 2022;43(10):1216-1221. doi:10.1097/MAO.0000000000003700

Summary of April 2023 Topic: The Vestibulo-Perceptual System

The first article investigated the vestibular perceptual threshold, physiological vestibular reflexes, and the development of motion sickness in patients with persistent postural-perceptual dizziness (PPPD). This cross-sectional study included 26 females diagnosed with PPPD and 33 age and gender-matched healthy controls (HC). All participants sat in a motorized rotary chair with no visual or auditory input for 20 minutes with slowly increasing velocity, up to a maximum of 72 degrees/second. Results showed lower vestibulo-perceptual thresholds and increased vegetative response (severe dizziness, nausea, or vomiting) in PPPD patients that were proportional to the duration of the disease. The authors concluded

Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

that the increased vestibular excitability likely reflects motion intolerance and the reproduction of dizziness when exposed to movement in PPPD patients (1).

The next article evaluated the function of the velocity-storage circuit by measuring postural and perceptual responses to a false inertial cue. Twenty healthy participants were taken through two different sessions, the control and test session, in which they were rotated clockwise, then counterclockwise, and with a different starting head position (head-down and head-up). The starting head position was maintained throughout the entire control session but was changed (at the end of rotation) during the test session. A significant correlation between postural and perceptual responses was shown as the head position and inertial motion perception at the end of the rotation matched with the inertia direction anticipated by the velocity-storage model. From this, the authors concluded that the postural instability clinically observed in vestibular disorders is likely from the pathomechanism of the velocity-storage circuit and its vestibulopostural response to a false vestibular cue (2).

The third article examined whether visual information can access short-latency vestibulo-spinal (collic) reflex. Twenty healthy participants were asked to rate the amount of self-motion (vection) they perceived while they viewed a large-field visual motion stimulation; cervical vestibular myogenic-evoked potentials (VEMPs) were simultaneously assessed. Results showed a positive correlation in the increased amplitude of the VEMPs with subjective vection strength. As a comparison, a small-field visual motion stimulation (nullifying vection perception) was done in a control group that failed to increase VEMP amplitude. Therefore, vection illusion can influence fast-activating reflexes, like VEMPs, and demonstrates the functional importance of perceptual mechanisms in human postural control (3).

The final article describes the importance of vestibular precision and presents findings on how precision compares across perception, the vestibulo-ocular reflex, posture, and neurons. The authors summarize how precision is an individual trait and that it correlates with clinical outcomes in patients and performance in functional tasks. The article concludes that the importance of continued vestibular precision and accuracy research can potentially improve outcomes in vestibular health and pathology (4).

1. Wurthmann, S., Holle, D., Obermann, M., Roesner, M., Nsaka, M., Scheffler, A., Kleinschnitz, C., & Naegel, S. (2021). Reduced vestibular perception thresholds in persistent postural-perceptual dizziness- a cross-sectional study. *BMC neurology*, 21(1), 394. <https://doi.org/10.1186/s12883-021-02417-z>
2. Choi, J. Y., Koo, Y. J., Song, J. M., Kim, H. J., & Kim, J. S. (2023). Effect of a False Inertial Cue in the Velocity-Storage Circuit on Head Posture and Inertia Perception. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 43(9), 1530–1539. <https://doi.org/10.1523/JNEUROSCI.1148-22.2023>
3. Bonsu, A. N., Nousi, S., Lobo, R., Strutton, P. H., Arshad, Q., & Bronstein, A. M. (2021). Vestibulo-perceptual influences upon the vestibulo-spinal reflex. *Experimental brain research*, 239(7), 2141–2149. <https://doi.org/10.1007/s00221-021-06123-7>

Summary of Monthly Abstract of the Week Topics From July 2017 to present

4. Diaz-Artiles, A., & Karmali, F. (2021). Vestibular Precision at the Level of Perception, Eye Movements, Posture, and Neurons. *Neuroscience*, 468, 282–320.
<https://doi.org/10.1016/j.neuroscience.2021.05.028>

Summary of May 2023 Topic: Meniere's

The May topic for abstract of the month was the topic of Meniere's. The first article looked at demographic data from a sample of patients with Meniere's disease. 520 patients were included where 81% of cases were unilateral. A positive family history of Meniere's was reported in 13% of participants, while a history of allergic diseases was reported in 33%. Further studies are still needed to examine more demographics around Meniere's disease. (1)

The second article looked to compare the different methods of audio-vestibular and contrast-enhanced MRI of inner ear in identifying endolymphatic hydrops which are key characteristics of Meniere's disease. There were 70 patients with unilateral Meniere's disease where audio-vestibular evaluations were performed, and the relationship between imaging signs of endolymphatic hydrops and audio-vestibular results was investigated. It was found that for the diagnosis of Meniere's disease that contrast-enhanced MRI was more advantageous in finding endolymphatic hydrops over the audio-vestibular evaluations. (2)

The last article looked to evaluate the pros and cons of intratympanic aminoglycosides versus either placebo or no treatment in people with Meniere's disease. There were 5 RCTs included with a total of 137 participants, and there were different outcomes looked at throughout the studies. It was found that there was an improvement in vertigo symptoms and global scores of vertigo were lower by participants who received the aminoglycoside treatment compared to no participants had an improvement in vertigo symptoms with the patients who received no treatment. It was also found that vertigo frequency was lower for those who received the aminoglycoside treatment compared to ones that received the placebo; these participants had an increase of vertigo frequency. From the studies, there were no risks found with usage of the treatment. But from all the studies, there were low participants, and different studies used different methods, so it was hard to determine if there was true clinical significance. (3)

1. Mammarella F, Loperfido A, Keeling EG, Bellocchi G, Marsili L. Ménière's Disease: Insights from an Italian Nationwide Survey. *Audiol Res*. 2023;13(2):160-168. Published 2023 Feb 28. doi:10.3390/audiolres13020016
2. Leng Y, Fan W, Liu Y, et al. Comparison between audio-vestibular findings and contrast-enhanced MRI of inner ear in patients with unilateral Ménière's disease. *Front Neurosci*. 2023;17:1128942. Published 2023 Mar 13. doi:10.3389/fnins.2023.1128942
3. Webster KE, Galbraith K, Lee A, et al. Intratympanic gentamicin for Ménière's disease. *Cochrane Database Syst Rev*. 2023;2(2):CD015246. Published 2023 Feb 27. doi:10.1002/14651858.CD015246.pub2

Summary of Monthly Abstract of the Week Topics From July 2017 to present

Summary of June 2023 Topic: Vascular Vertigo

The June topic for the abstract of the month was Vascular Vertigo. The first article presented the diagnostic criteria for vascular vertigo and dizziness based on the Barany Society. The article includes diagnostic criteria for acute prolonged and transient vascular vertigo/dizziness, Vertebral Artery Compression Syndrome, and lesion sites responsible for isolated vascular vertigo. Vascular vertigo/dizziness should be considered in patients with acute vestibular symptoms that show findings of central HINTS (normal head-impulse test, direction-changing gaze-evoked nystagmus, or pronounced skew deviation) and have vascular risk factors. The authors conclude that identifying the underlying etiology of vascular vertigo/dizziness can help guide appropriate medical management options to treat acute stroke and secondary prevention of future stroke (1).

The next article described the clinical symptoms and laboratory vestibular findings of Vestibular Paroxysmia (VP) and its long-term clinical course. Based on the Barany Society diagnostic criteria, 146 patients with VP were identified. The clinical symptoms found were: frequency (5 to 30 attacks per day); the most common symptom was spinning vertigo; two-thirds of patients experienced spontaneous attacks; head movements triggered one-quarter of patients; 70% of patients reported no accompanying symptoms. Vestibular findings were as follows: one-third of patients showed hyperventilation-induced nystagmus; 3 to 6 degrees of deviation in subjective visual vertical; and no complete loss of peripheral vestibular function was evident. The long-term course of VP shows a reduction in attack frequency in which most patients did not require medication or ongoing treatment (2).

The third article summarized the evidence and the accuracy of neuro-otologic bedside examination techniques in differentiating central versus peripheral causes of acute vestibular syndrome (AVS). This systematic review and meta-analysis included 14 articles for quantitative analysis reporting on the diagnostic accuracy of physical examination in adults with acute, prolonged dizziness/vertigo. Results revealed that the Head Impulse, Nystagmus, Test of Skew (HINTS) had 95.3% sensitivity and 92.6% specificity, and early (within 24-48 hours) MRI-diffusion-weighted imaging (DWI) was falsely negative in 15% of strokes. The authors concluded that the HINTS examination, performed by appropriately trained clinicians, has higher diagnostic accuracy in identifying stroke than early MRI-DWI and can help determine central from peripheral causes in AVS (3).

The final article presents a case report of a patient with a rare subtype of infarction to the medial medulla oblongata, Spiller syndrome. The 42-year-old patient with a history of hypertension, diabetes mellitus, and cerebral hemorrhage initially showed signs and symptoms consistent with acute vestibular neuritis. However, symptoms progressed to left-sided limb weakness and dysarthria after 4 hours of admission to the emergency department. Bedside exam revealed ipsilateral hypoglossal paralysis and contralateral loss of deep sensation. MRI confirmed medial medullary infarction (MMI). Authors highlight the importance of considering a central lesion in patients with acute vestibular syndrome with vascular risk factors (4).

1. Kim, J. S., Newman-Toker, D. E., Kerber, K. A., Jahn, K., Bertholon, P., Waterston, J., & Strupp, M. (2022). Vascular vertigo and dizziness: Diagnostic criteria. *Journal of Vestibular Research*, (Preprint), 1-18.

Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

2. Steinmetz, K., Becker-Bense, S., Strobl, R., Grill, E., Seelos, K., & Huppert, D. (2022). Vestibular paroxysmia: clinical characteristics and long-term course. *Journal of Neurology*, 269(12), 6237-6245.
3. Tarnutzer, A. A., Gold, D., Wang, Z., Robinson, K. A., Kattah, J. C., Mantokoudis, G., ... & Newman-Toker, D. E. (2023). Impact of clinician training background and stroke location on bedside diagnostic accuracy in the acute vestibular syndrome—a meta-analysis. *Annals of neurology*.
4. Wang, H., Shi, T., Shang, Y., Chen, X., Xu, J., & Geng, Y. (2022). Case report: Spiller syndrome initially mimicking vestibular neuritis. *Frontiers in Neurology*, 13.

Summary of July 2023 Topic: Persistent Postural-Perceptual Dizziness (PPPD)

The July topic for abstract of the month was 3PD. The first article looked at pharmacological interventions for 3PD and analyzed the benefits vs harms of different interventions. There was a systematic review done that found no evidence from placebo controlled randomized trials regarding treatment. Therefore, there is big uncertainty over the use of these treatments, and more research is needed to decide if there are effective treatments and if there are any adverse effects of the treatment.

(1)

The second article looked to examine the effect on gaze instability after exposure to moving or complex visual stimuli in patients with 3PD. It was found that patients with 3PD were more likely to exhibit gaze instability after exposure to moving visual stimuli which caused an exacerbation of vestibular symptoms.

(2)

The next article looked at the intervention of cognitive- behavioral therapy in treating 3PD compared to the gold standard of physical therapy. 40 adults with 3PD were randomized to 6 sessions of the cognitive behavioral therapy or to the physical therapy treatment group. Results showed that there were small to moderate treatment effects in favor of the cognitive behavioral therapy.

(3)

The last article looked to compare the results of specific outcome measures in patients with vestibular migraines versus 3PD. 30 patients were put into 3 groups: vestibular migraine, 3PD, and healthy volunteers where there were multiple outcome measures assessed. It was found that Dizziness Handicap Inventory, anxiety and somatic symptom scores, and depression scores was significantly high in both the migraine and 3PD group. Emotional and functional scores, anxiety and somatization, and mental health and social role were worse for the 3PD patients.

(4)

1. Webster KE, Harrington-Benton NA, Judd O, Kaski D, Maarsingh OR, MacKeith S, Ray J, Van Vugt VA, Burton MJ. Pharmacological interventions for persistent postural-perceptual dizziness (PPPD). *Cochrane Database Syst Rev*. 2023 Mar 9;3(3):CD015188. doi: 10.1002/14651858.CD015188.pub2. PMID: 36906836; PMCID: PMC9997546.

Summary of Monthly Abstract of the Week Topics From July 2017 to present

2. Yagi C, Morita Y, Yamagishi T, Ohshima S, Izumi S, Takahashi K, Itoh K, Suzuki Y, Igarashi H, Horii A. Gaze instability after exposure to moving visual stimuli in patients with persistent postural-perceptual dizziness. *Front Hum Neurosci.* 2022 Nov 25;16:1056556. doi: 10.3389/fnhum.2022.1056556. PMID: 36504627; PMCID: PMC9733075.
3. Herdman D, Norton S, Murdin L, Frost K, Pavlou M, Moss-Morris R. The INVEST trial: a randomised feasibility trial of psychologically informed vestibular rehabilitation versus current gold standard physiotherapy for people with Persistent Postural Perceptual Dizziness. *J Neurol.* 2022 Sep;269(9):4753-4763. doi: 10.1007/s00415-022-11107-w. Epub 2022 Apr 10. PMID: 35397754; PMCID: PMC8994825.
4. Ak AK, Çelebisoy N, Özdemir HN, Gökçay F. Vestibular migraine and persistent postural perceptual dizziness: Handicap, emotional comorbidities, quality of life and personality traits. *Clin Neurol Neurosurg.* 2022 Oct;221:107409. doi: 10.1016/j.clineuro.2022.107409. Epub 2022 Aug 5. PMID: 35961229.

Summary of August 2023 Topic: Expectancies and Predictive Processing

The first article aimed to explain the new empirical concept of Brainstem Cortical Scaling Metric (BCSM) and its primary role in vestibulo-spatial and vestibulo-temporal perceptual maps of the external world with its associated action-perceptual risk assessment. The authors provided empirical data for this concept using two techniques (combining binocular rivalry with vestibular stimulation and modulating vestibulo-cortical excitability over the posterior parietal cortex) to disrupt interhemispheric interactions that may play a role in balance control. They found and concluded that using the BCSM concept, time and space were distorted, which impairs the accurate assessment of errors between predicted and actual motion and, therefore, mechanistically accounts for functional dizziness (1).

The next article investigated the relationship between objective sway and subjective instability by modulating prior expectations regarding the associated difficulty of an upcoming postural task via priming in two groups. The visual priming group consisted of thirty healthy participants who watched a video of a stable or unstable actor performing a postural task. The verbal priming group consisted of thirty healthy individuals who were either given verbal erroneous information regarding how much the platform would be moving or no information. In the visual group, those who watched the unstable actor did not maintain a normal relationship between trunk sway and subjective instability compared to the stable actor, and the verbal priming group showed an increase in subjective instability in those who were told the platform amplitude would increase, even though it did not. Therefore, the authors concluded that patients with balance disorders may benefit from enhanced cognitive retraining, primarily modifying expectations (2).

The third article provided literature on how predictive processing concepts can be applied to the interaction of motor control with anxiety. The Predictive Processing Framework (PPF) views the central nervous system as a hierarchical prediction engine that determines top-down hypotheses about the

Summary of Monthly Abstract of the Week Topics From July 2017 to present

state of the body and the world and minimizes prediction error over time. Many examples were presented using predictive processing concepts such as attention, precision, and motor control to explain the fear of falling in older adults and “choking” in athletes. The authors conclude that there is potential for predictive processing to guide future work in generating and testing new hypotheses in anxiety-induced disruptions to motor control and support the design of interventions (3).

The fourth article investigated the role of cognitive factors and personality traits on the perception of vection. Fifty-one healthy participants were divided into three groups based on given vection sensation expectations: strong expectation, weak expectation, or control group. The configuration of the chair (either rotatable or fixed) was also modified during the experiment. Results revealed that expectation directly affected perceived vection intensity but was limited to the rotatable chair. In addition, field dependence and depersonalization traits were found to have a positive correlation with vection perception. The authors concluded that vection should be considered more than a perceptual phenomenon as this study showed the impact of high-level cognitive factors (e.g., expectation) and personality traits on vection perception (4).

The final article presented recent research about the proprioceptive and vestibular systems interaction and the neural computations underlying the predictive sense of active self-motion. It summarized different data and articles that discuss the suppression of vestibular inputs during active head movements, expected and actual proprioceptive feedback during self-motion, the proprioceptive system during active and passive movements, vestibular-propriceptive integration, and proprioceptive substitution in vestibular pathways following vestibular sensory loss (5).

1. Arshad, Q., Saman, Y., Sharif, M., Kaski, D., & Staab, J. P. (2022). Magnitude Estimates Orchestrate Hierarchical Construction of Context-Dependent Representational Maps for Vestibular Space and Time: Theoretical Implications for Functional Dizziness. *Frontiers in integrative neuroscience*, 15, 806940. <https://doi.org/10.3389/fnint.2021.806940>
2. Castro, P., Papoutselou, E., Mahmoud, S., Hussain, S., Bassaletti, C. F., Kaski, D., Bronstein, A., & Arshad, Q. (2022). Priming overconfidence in belief systems reveals negative return on postural control mechanisms. *Gait & posture*, 94, 1–8. <https://doi.org/10.1016/j.gaitpost.2022.02.015>
3. Harris, D. J., Wilkinson, S., & Ellmers, T. J. (2023). From fear of falling to choking under pressure: A predictive processing perspective of disrupted motor control under anxiety. *Neuroscience and biobehavioral reviews*, 148, 105115. <https://doi.org/10.1016/j.neubiorev.2023.105115>
4. D'Amour, S., Harris, L. R., Berti, S., & Keshavarz, B. (2021). The role of cognitive factors and personality traits in the perception of illusory self-motion (vection). *Attention, perception & psychophysics*, 83(4), 1804–1817. <https://doi.org/10.3758/s13414-020-02228-3>
5. Cullen, K. E., & Zobeiri, O. A. (2021). Proprioception and the predictive sensing of active self-motion. *Current opinion in physiology*, 20, 29–38. <https://doi.org/10.1016/j.cophys.2020.12.001>

Summary of Monthly Abstract of the Week Topics From July 2017 to present

Summary of September 2023 Topic: BPPV Updates

The September topic for abstract of the month was BPPV updates. The first article looked at a new treatment for treating canalolithiasis-type BPPV. There was a 3D model of the human labyrinth that was used to analyze the maneuver. The results showed that the 3D model predicted the effectiveness of the maneuver for the treatment of either single canal or multiple canal BPPV canalolithiasis, making treatment more straightforward. (1)

The next article looked to evaluate the effectiveness of the head rotation test with bowing in the sitting position in diagnosing lateral semicircular canal BPPV. 25 patients with lateral semicircular canal BPPV were evaluated using the head rotation test with bowing, and the traditional head rotation in the supine position and the bow and lean test were assessed to compare. It was found that the head rotation test with bowing identified direction changing nystagmus in lateral semicircular canal BPPV, and it was found that it could pinpoint the determination of the affected ears. (2)

The next article looked to assess gait, falls, and fear of falling because of having BPPV and being treated with the particle repositioning maneuvers. From the review, it was found that people with BPPV walked slower in general and walked slower during head rotations and demonstrated more sway during tandem walking compared with the controls. It was found that after the particle repositioning maneuver that gait speed increased and became safer. Difficulties during tandem walking and walking with head rotations did not improve. Falls were increasingly higher for patients with BPPV than for controls, but after treatment, the number of falls and fear of falling decreased. To conclude, BPPV increases the chances of falls and impacts the spatiotemporal parameters of GAIT. (3)

The last article evaluated the extent to which BPPV is associated with depression and anxiety. It was found that BPPV was associated with an increased risk of anxiety compared to controls, but there is still more research needed for more confirmation. (4)

1. Barreto RG, Yacovino DA, Cherchi M, Teixeira LJ, Nader SN, Leão GF. Universal Repositioning Maneuver: A New Treatment for Single Canal and Multi-Canal Benign Paroxysmal Positional Vertigo by 3-Dimensional Model Analysis. *J Int Adv Otol.* 2023;19(3):242-247. doi:10.5152/iao.2023.22921
2. Kim MW, Jin MH, Kim JK, Kwak MY. Efficacy of the Head Rotation Test With Bowing for the Lateral Canal Benign Paroxysmal Positional vertigo [published online ahead of print, 2023 Aug 9]. *Otol Neurotol.* 2023;10.1097/MAO.0000000000003982. doi:10.1097/MAO.0000000000003982
3. Pauwels S, Casters L, Lemkens N, et al. Gait and Falls in Benign Paroxysmal Positional Vertigo: A Systematic Review and Meta-analysis. *J Neurol Phys Ther.* 2023;47(3):127-138. doi:10.1097/NPT.0000000000000438
4. Yeo BSY, Toh EMS, Lim NE, et al. Association of Benign Paroxysmal Positional Vertigo with Depression and Anxiety-A Systematic Review and Meta-Analysis [published online ahead of print, 2023 Aug 10]. *Laryngoscope.* 2023;10.1002/lary.30957. doi:10.1002/lary.30957



Vestibular Rehabilitation SIG

Summary of Monthly Abstract of the Week Topics From July 2017 to present

Summary of October 2023 Topic: Aging and Vestibular Dysfunction

Summary of November 2023 Topic:

Summary of December 2023 Topic: